arXiv:1009.3370v3 [math.RT] 16 Jul 2011

SILTING MUTATION IN TRIANGULATED CATEGORIES

TAKUMA AIHARA AND OSAMU IYAMA

Abstract. In representation theory of algebras the notion of 'mutation' often plays important roles, and two cases are well known, i.e. 'cluster tilting mutation' and 'exceptional mutation'. In this paper we focus on 'tilting mutation', which has a disadvantage that it is often impossible, i.e. some of summands of a tilting object can not be replaced to get a new tilting object. The aim of this paper is to take away this disadvantage by introducing 'silting mutation' for silting objects as a generalization of 'tilting mutation'. We shall develop a basic theory of silting mutation. In particular, we introduce a partial order on the set of silting objects and establish the relationship with 'silting mutation' by generalizing the theory of Riedtmann-Schofield and Happel-Unger. We show that iterated silting mutation act transitively on the set of silting objects for local, hereditary or canonical algebras. Finally we give a bijection between silting subcategories and certain t-structures.

Contents

1. Introduction 1
2. Silting subcategories 4
2.1. Definition and basic properties 4
2.2. Partial order on silting subcategories 5
2.3. Krull-Schmidt triangulated categories 7
2.4. Silting mutation 10
2.5. Silting reduction 13
2.6. Silting quivers and examples 16
2.7. Okuyama-Rickard complexes and APR and BB tilting modules 19
3. Transitivity for piecewise hereditary algebras 21
4. Silting subcategories in triangulated categories with coproducts 24
4.1. Torsion pairs induced by sets of compact objects 24
4.2. The correspondence between silting subcategories and t-structures 27
References 28

1. Introduction

In representation theory of algebras the notion of 'mutation' often plays important roles. Mutation is an operation for a certain class of objects (e.g. cluster tilting objects) in a fixed category to construct a new object from a given one by replacing a summand. Two important cases are well-known: One is 'cluster tilting mutation' [BMRRT, IY, IW] for cluster tilting objects which is applied for categorification of Fomin-Zelevinsky cluster algebras, and the other is 'exceptional mutation' for exceptional sequences which is used to study the structure of derived categories of algebraic varieties [GR, C].

From Morita theoretic viewpoint [Ric1], tilting objects are the most important class of objects, and 'tilting mutation' for tilting objects has been studied by several authors. 'Tilting mutation' has the origin in BGP (=Bernstein-Gelfand-Ponomarev) reflection functor [BGP] in quiver representation theory, and reflection functors are understood as a special class of APR (=Auslander-Platzeck-Reiten) tilting modules [APR], which are 'tilting mutations' obtained by replacing a simple summand of the tilting A-modules A. In 1991 general 'tilting mutation' for tilting modules was introduced by Riedtmann-Schofield [RS] in their

2010 *Mathematics Subject Classification.* 16E30, 18E30.

study of combinatorial aspects of tilting theory. It has been shown by Happel-Unger [HU] that 'tilting mutation' is closely related to the partial order of tilting modules given by the inclusion relation of the associated t-structures, and this is a big advantage of 'tilting mutation' which 'cluster tilting mutation' and 'exceptional mutation' do not have. Their beautiful theory of 'tilting mutation' has a lot of important applications, especially in the study of cluster categories and preprojective algebras [BMRRT, IR, BIRS, BIKR, SY]. Another important source of 'tilting mutation' is modular representation theory, where 'tilting mutation' for tilting complexes was introduced by Okuyama and Rickard [O, Ric2, HK], and has played an important role in the study of Broué's abelian defect group conjecture.

It is remarkable that 'tilting mutation' has a big disadvantage that it is often impossible, i.e. some of summands of a tilting object can not be replaced to get a new tilting object. So it is usual that we can not get sufficiently many tilting objects by iterated 'tilting mutation'. For example the tilting A-module DA usually can not be obtained by iterated 'tilting mutation' of A even for the case of hereditary algebras A. The aim of this paper is to take away this disadvantage by introducing '*silting mutation*' for *silting objects*. In this context 'tilting mutation' should be understood as a special case of 'silting mutation'. Actually classical notion of APR and BB tilting modules and Okuyama-Rickard complexes can be understood as special cases of 'silting mutation' (Theorems 2.53 and 2.50). Silting objects are generalization of tilting objects, and sometimes appeared in representation theory mainly in the study of t-structures, e.g. Keller-Vossieck [KV], Hoshino-Kato-Miyachi [HKM], Assem-Salorio-Trepode [AST] and Wei [W]. A point is that 'silting mutation' is always possible in the sense that any summand of a silting object always can be replaced to get a new silting object (Theorem 2.31). Hence it is natural to hope that 'silting mutation' gives us sufficiently many silting objects in triangulated categories. We pose the following question (Question 2.42), where the corresponding property for tilting objects is usually not satisfied.

Question 1.1. Let A be a finite dimensional algebra over a field. When does A satisfy the following property (T) (respectively, (T′))?

(T) (respectively, (T′)) The action of iterated irreducible 'silting mutation' (respectively, iterated 'silting mutation') on the set of basic silting objects in $\mathsf{K}^{\mathrm{b}}(\text{proj-}A)$ is transitive.

We shall show the following partial answers in Corollary 2.43 and Theorem 3.1.

Theorem 1.2. *If A is either a local algebra, a hereditary algebra or a canonical algebra, then (T) is satisfied.*

It will be shown in [A] that (T) is satisfied also for representation-finite symmetric algebras. On the other hand, there is a symmetric algebra such that (T) is not satisfied [AGI]. We do not know any algebra such that (T′) is not satisfied.

As a basic tool of the study of silting objects, we shall introduce a partial order on silting objects (Theorem 2.11) and establish the relationship with 'silting mutation' (Theorem 2.35) by generalizing the theory of Riedtmann-Schofield and Happel-Unger for tilting modules. In particular, Question 1.1 is equivalent to the following question.

Question 1.3. Let A be a finite dimensional algebra over a field. When is the Hasse quiver of the partially ordered set of basic silting objects in $\mathsf{K}^{\mathrm{b}}(\text{proj-}A)$ connected?

The comparison of three kinds of mutation is explained by the following table.

	always possible	partial order exists	transitive for hereditary case
cluster tilting mutation	Yes	No	Yes
silting mutation	Yes	Yes	Yes
∪			
tilting mutation	No	Yes	No
exceptional mutation	Yes	No	Yes

In our paper we study basic properties of silting objects/subcategories. We show that non-isomorphic indecomposable objects in any silting subcategory form a basis of the Grothendieck group of the triangulated category (Theorem 2.27). In particular all basic silting objects have the same number of indecomposable summands (Corollary 2.28). We also introduce 'silting reduction' (Theorem 2.37), which gives a bijection between silting subcategories containing a certain fixed subcategory and silting subcategories in a certain quotient triangulated category.

In section 4.2 we study in detail the relationship between silting subcategories and t-structures under the assumption that the triangulated categories have arbitrary coproducts, and improve some of pioneering results of Hoshino-Kato-Miyachi [HKM]. The advantage of this setting is that each set of compact objects gives rise to a torsion pair (Theorem 4.3), which is not the case for the setting of section 2. This basic result seems to be new and to have independent interest. As an application, we establish a one-to-one correspondence between silting subcategories and t-structures satisfying certain conditions (Theorem 4.10).

We notice that a different generalization of 'tilting mutation' was given in [IO, HX]. Also some aspects of mutation were recently discussed in [BRT, L, KY].

Parts of results in this paper were presented in Trondheim (March 2009), Nagoya (June 2009), Matsumoto (October 2009), Shizuoka (November 2009) and Tokyo (August 2010).

Notations Let $\mathcal{T}$ be an additive category. For morphisms $f : X \to Y$ and $g : Y \to Z$ in a category, we denote by $gf : X \to Z$ the composition. We say that a morphism $f : X \to Y$ is *right minimal* if any morphism $g : X \to X$ satisfying $fg = f$ is an isomorphism. Dually we define a *left minimal* morphism.

For a collection $\mathcal{X}$ of objects in $\mathcal{T}$, we denote by $\mathsf{add}\mathcal{X}$ (respectively, $\mathsf{Add}\mathcal{X}$) the smallest full subcategory of $\mathcal{T}$ which is closed under finite (respectively, arbitrary) coproducts, summands and isomorphisms and contains $\mathcal{X}$. We denote by $\mathsf{smd}\mathcal{X}$ the smallest full subcategory of $\mathcal{T}$ which is closed under summands and contains $\mathcal{X}$. When we say that $\mathcal{X}$ is a *subcategory* of $\mathcal{T}$, we always assume that $\mathcal{X}$ is full and satisfies $\mathcal{X} = \mathsf{add}\mathcal{X}$.

Let $\mathcal{X}$ be a subcategory of $\mathcal{T}$. We say that a morphism $f : X \to Y$ is a *right $\mathcal{X}$-approximation* of Y if $X \in \mathcal{X}$ and $\mathrm{Hom}_{\mathcal{T}}(X, f)$ is surjective for any $X \in \mathcal{X}$. We say that $\mathcal{X}$ is *contravariantly finite* if any object in $\mathcal{T}$ has a right $\mathcal{X}$-approximation. Dually, we define a *left $\mathcal{X}$-approximation* and a *covariantly finite subcategory*. We say that $\mathcal{X}$ is *functorially finite* if it is contravariantly and covariantly finite.

When $\mathcal{T}$ is a triangulated category, we denote by $\mathsf{thick}\mathcal{X}$ the smallest thick subcategory of $\mathcal{T}$ containing $\mathcal{X}$. For collections $\mathcal{X}$ and $\mathcal{Y}$ of objects in $\mathcal{T}$, we denote by $\mathcal{X} * \mathcal{Y}$ the collection of objects $Z \in \mathcal{T}$ appearing in a triangle $X \to Z \to Y \to X[1]$ with $X \in \mathcal{X}$ and $Y \in \mathcal{Y}$. We set

$$\begin{array}{rcl} \mathcal{X}^{\perp} = \mathcal{X}^{\perp_{\mathcal{T}}} & := & \{T \in \mathcal{T} \mid \mathrm{Hom}_{\mathcal{T}}(\mathcal{X}, T) = 0\}, \\ {}^{\perp}\mathcal{X} = {}^{\perp_{\mathcal{T}}}\mathcal{X} & := & \{T \in \mathcal{T} \mid \mathrm{Hom}_{\mathcal{T}}(T, \mathcal{X}) = 0\} \end{array}$$

For an additive category $\mathcal{A}$, we denote by $\mathsf{K}^{\mathrm{b}}(\mathcal{A})$ the homotopy category of bounded complexes over $\mathcal{A}$. For an abelian category $\mathcal{A}$, we denote by $\mathsf{D}^{\mathrm{b}}(\mathcal{A})$ the bounded derived category of $\mathcal{A}$.

For a ring A, we denote by Mod-A (respectively, mod-A) the category of all (respectively, finitely generated) right A-modules, by proj-A the category of finitely generated projective A-modules. We denote by J_A the Jacobson radical of A. When A is a finite dimensional algebra over a field k, we denote by $D := \mathrm{Hom}_k(-, k) : \text{mod-}A \leftrightarrow \text{mod-}A^{\mathrm{op}}$ the k-duality.

Definition 1.4. Let $\mathcal{T}$ be a triangulated category and $(\mathcal{X}, \mathcal{Y})$ a pair of subcategories of $\mathcal{T}$.

- We say that $(\mathcal{X}, \mathcal{Y})$ is a *torsion pair* if $\mathrm{Hom}_{\mathcal{T}}(\mathcal{X}, \mathcal{Y}) = 0$ and $\mathcal{X} * \mathcal{Y} = \mathcal{T}$.
- We say that $(\mathcal{X}, \mathcal{Y})$ is a *t-structure* [BBD] if $(\mathcal{X}[1], \mathcal{Y})$ is a torsion pair and $\mathcal{X}[1] \subset \mathcal{X}$. In this case $\mathcal{X} \cap \mathcal{Y}$ is called the *heart*.
- We say that $(\mathcal{X}, \mathcal{Y})$ is a *co-t-structure* [P1] if $(\mathcal{X}[-1], \mathcal{Y})$ is a torsion pair and $\mathcal{X} \subset \mathcal{X}[1]$. In this case $\mathcal{X} \cap \mathcal{Y}$ is called the *coheart*.
- We say that a torsion pair $(\mathcal{X}, \mathcal{Y})$ is a *stable t-structure* [M] if $\mathcal{X} = \mathcal{X}[1]$.

Acknowledgements The second author would like to thank Jiaqun Wei for stimulating discussion. The authors would like to thank David Pauksztello, Apostolos Beligiannis and Sonia Trepode for useful information.

2. Silting subcategories

Let $\mathcal{T}$ be a triangulated category. We do not assume anything else on $\mathcal{T}$ unless otherwise stated. Throughout this paper we write the vanishing condition $\mathrm{Hom}_{\mathcal{T}}(X, Y[i]) = 0$ for any $i \in \mathbb{Z}$ by

$$\mathrm{Hom}_{\mathcal{T}}(X, Y[\mathbb{Z}]) = 0$$

simply. Similarly we often use

$$\mathrm{Hom}_{\mathcal{T}}(X, Y[>0]) = 0,\ \mathrm{Hom}_{\mathcal{T}}(X, Y[<0]) = 0,\ \mathrm{Hom}_{\mathcal{T}}(X, Y[\neq 0]) = 0 \ \text{ and so on.}$$

2.1. **Definition and basic properties.** In this subsection we introduce silting subcategories/objects of triangulated categories and study their basic properties.

Definition 2.1. Let $\mathcal{M}$ be a subcategory of a triangulated category $\mathcal{T}$.

(a) We say that $\mathcal{M}$ is *silting* if $\mathrm{Hom}_{\mathcal{T}}(\mathcal{M}, \mathcal{M}[>0]) = 0$ and $\mathcal{T} = \mathsf{thick}\mathcal{M}$. We denote by $\mathrm{silt}\,\mathcal{T}$ the collection of silting subcategories in $\mathcal{T}$.
(b) We say that $\mathcal{M}$ is *tilting* if $\mathrm{Hom}_{\mathcal{T}}(\mathcal{M}, \mathcal{M}[\neq 0]) = 0$ and $\mathcal{T} = \mathsf{thick}\mathcal{M}$.
(c) We say that an object $M \in \mathcal{T}$ is *silting* (respectively, *tilting*) if so is $\mathsf{add}M$.

The following examples give typical classes of tilting/silting objects.

Example 2.2. (a) Let A be a ring. Then A regarded as a stalk complex is a tilting object in $\mathsf{K}^{\mathrm{b}}(\text{proj-}A)$. More generally, any tilting complex of A is a tilting object in $\mathsf{K}^{\mathrm{b}}(\text{proj-}A)$.
(b) Let A be a differential graded ring and let $\mathsf{D}(A)$ be the derived category of A. If $H^i(A) = 0$ for any $i > 0$, then the thick subcategory $\mathsf{thick}(A)$ of $\mathsf{D}(A)$ generated by A has a silting object A.

The above example (a) is standard among triangulated categories with tilting objects in the following sense:

Proposition 2.3. [K] *Let $\mathcal{T}$ be an algebraic triangulated category. If $\mathcal{T}$ has a tilting object M, then $\mathcal{T}$ is triangle equivalent to* $\mathsf{K}^{\mathrm{b}}(\text{proj-}\mathrm{End}_{\mathcal{T}}(M))$.

On the other hand, we have the following necessary conditions for existence of silting/tilting subcategories.

Proposition 2.4. *Let $\mathcal{T}$ be a triangulated category. If $\mathcal{T}$ contains a silting (respectively, tilting) subcategory, then for any $X, Y \in \mathcal{T}$ we have* $\mathrm{Hom}_{\mathcal{T}}(X, Y[i]) = 0$ *for $i \gg 0$ (respectively, $|i| \gg 0$).*

Proof. We only show the statement for silting subcategories. Let $\mathcal{U} := \{X \in \mathcal{T} \mid \mathrm{Hom}_{\mathcal{T}}(X, M[\gg 0]) = 0 \text{ for any } M \in \mathcal{M}\}$. Then $\mathcal{U}$ is a thick subcategory of $\mathcal{T}$ containing $\mathcal{M}$. Thus we have $\mathcal{U} = \mathcal{T}$ since $\mathsf{thick}\mathcal{M} = \mathcal{T}$. For any $X \in \mathcal{T}$, let $\mathcal{V}_X := \{Y \in \mathcal{T} \mid \mathrm{Hom}_{\mathcal{T}}(X, Y[\gg 0]) = 0\}$. Then we have $\mathcal{M} \subset \mathcal{V}_X$ by the above argument. Since $\mathcal{V}_X$ is a thick subcategory of $\mathcal{T}$ containing $\mathcal{M}$, we have $\mathcal{V}_X = \mathcal{T}$ again. Thus the assertion holds. □

Immediately we have the following observation.

Example 2.5. Let A be a finite dimensional algebra over a field k.

(a) $\mathsf{D}^{\mathrm{b}}(\text{mod-}A)$ contains silting subcategories if and only if $\mathrm{gl.dim}A < \infty$.
(b) Assume that A is self-injective. Then the stable module category $\underline{\text{mod}}\text{-}A$ of mod-A contains silting subcategories if and only if A is semisimple.

Proof. We only have to show 'only if' part. Both for the cases (a) and (b), we have $\mathrm{Ext}^i_A(A/J_A, A/J_A) \neq 0$ for any $i > 0$. Thus the assertions follow from Proposition 2.4. □

It is often the case that triangulated categories have many silting subcategories which are not tilting. But there are important triangulated categories such that all silting subcategories are tilting.

Definition 2.6. Let ℓ be an integer. We call a k-linear Hom-finite triangulated category $\mathcal{T}$ *ℓ-Calabi-Yau* if there is a bifunctorial isomorphism $\mathrm{Hom}_{\mathcal{T}}(-, ?) \simeq D\mathrm{Hom}_{\mathcal{T}}(?, -[\ell])$.

One can easily check the following statements.

Lemma 2.7. *Let ℓ be an integer and $\mathcal{T}$ an ℓ-Calabi-Yau triangulated category. If $\mathcal{T} \neq 0$, then the following assertions hold:*
(1) *If $\ell = 0$, then any silting subcategory of $\mathcal{T}$ is tilting;*
(2) *If $\ell > 0$, then there exist no silting subcategories of $\mathcal{T}$;*
(3) *If $\ell < 0$, then there exist no tilting subcategories of $\mathcal{T}$.*

Proof. Since any non-zero object $X \in \mathcal{T}$ satisfies $\mathrm{Hom}_{\mathcal{T}}(X, X[\ell]) \simeq D\mathrm{Hom}_{\mathcal{T}}(X, X) \neq 0$, the assertions (2) and (3) follow. We can prove (1) similarly. □

Immediately we have the following observation.

Example 2.8. Let A be a finite dimensional symmetric algebra over a field. Then any silting object in $\mathsf{K}^{\mathrm{b}}(\text{proj-}A)$ is tilting.

Proof. We know that $\mathsf{K}^{\mathrm{b}}(\text{proj-}A)$ is a 0-Calabi-Yau triangulated category by Auslander-Reiten duality [H]. Thus the assertion follows from Lemma 2.7. □

Notice that we do not know whether there exist ℓ-Calabi-Yau triangulated categories containing a silting subcategory and $\ell < 0$.

We end this subsection with the following remark, where we say that a category is *skeletally small* if isomorphism classes of objects form a set.

Remark 2.9. If a triangulated category $\mathcal{T}$ has a skeletally small silting subcategory, then $\mathcal{T}$ is also skeletally small (e.g. Proposition 2.17).

2.2. **Partial order on silting subcategories.** The aim of this subsection is to introduce a partial order on silting subcategories as a generalization of the partial order on tilting modules introduced by Riedtmann-Schofield [RS] and Happel-Unger [HU]. Our main result in this subsection is Theorem 2.11 below.

Definition 2.10. For $\mathcal{M}, \mathcal{N} \in \mathrm{silt}\,\mathcal{T}$, we write $\mathcal{M} \geq \mathcal{N}$ if $\mathrm{Hom}_{\mathcal{T}}(\mathcal{M}, \mathcal{N}[>0]) = 0$.

Theorem 2.11. *$\geq$ gives a partial order on* $\mathrm{silt}\,\mathcal{T}$.

The following subcategory plays an important role in the proof of Theorem 2.11.

Definition 2.12. For any $\mathcal{M} \in \mathrm{silt}\,\mathcal{T}$, we define a subcategory of $\mathcal{T}$ by

$$\mathcal{T}_{\mathcal{M}}^{\leq 0} := \{X \in \mathcal{T} \mid \mathrm{Hom}_{\mathcal{T}}(\mathcal{M}, X[>0]) = 0\}.$$

We put $\mathcal{T}_{\mathcal{M}}^{\leq \ell} = \mathcal{T}_{\mathcal{M}}^{<\ell+1} := \mathcal{T}_{\mathcal{M}}^{\leq 0}[-\ell]$ for $\ell \in \mathbb{Z}$.

Clearly we have $\mathcal{M} \subset \mathcal{T}_{\mathcal{M}}^{\leq 0}$ and $\mathcal{T}_{\mathcal{M}}^{<0} \subset \mathcal{T}_{\mathcal{M}}^{\leq 0}$. For any non-zero object $X \in \mathcal{T}$, we have $X[\gg 0] \in \mathcal{T}_{\mathcal{M}}^{\leq 0}$ and $X[\ll 0] \notin \mathcal{T}_{\mathcal{M}}^{\leq 0}$ by Proposition 2.4.

Crucial results are the following.

Proposition 2.13. *For any $\mathcal{M} \in \mathrm{silt}\,\mathcal{T}$, we have $\mathcal{T}_{\mathcal{M}}^{\leq 0} \cap {}^{\perp}(\mathcal{T}_{\mathcal{M}}^{<0}) = \mathcal{M}$.*

Proposition 2.14. *Let $\mathcal{M}, \mathcal{N} \in \mathrm{silt}\,\mathcal{T}$. Then $\mathcal{M} \geq \mathcal{N}$ if and only if $\mathcal{T}_{\mathcal{M}}^{\leq 0} \supset \mathcal{T}_{\mathcal{N}}^{\leq 0}$.*

Before proving these Propositions, we prove Theorem 2.11 by using them.

(i) We shall show that $\mathcal{M} \geq \mathcal{N}$ and $\mathcal{N} \geq \mathcal{L}$ imply $\mathcal{M} \geq \mathcal{L}$. By Proposition 2.14, we have $\mathcal{T}_{\mathcal{M}}^{\leq 0} \supset \mathcal{T}_{\mathcal{N}}^{\leq 0} \supset \mathcal{T}_{\mathcal{L}}^{\leq 0}$. Thus $\mathcal{T}_{\mathcal{M}}^{\leq 0} \supset \mathcal{T}_{\mathcal{L}}^{\leq 0}$, and we have $\mathcal{M} \geq \mathcal{L}$ again by Proposition 2.14.

(ii) We shall show that $\mathcal{M} \geq \mathcal{N}$ and $\mathcal{N} \geq \mathcal{M}$ imply $\mathcal{M} = \mathcal{N}$. By Proposition 2.14, we have $\mathcal{T}_{\mathcal{M}}^{\leq 0} = \mathcal{T}_{\mathcal{N}}^{\leq 0}$. Thus we have ${}^{\perp}(\mathcal{T}_{\mathcal{M}}^{<0}) = {}^{\perp}(\mathcal{T}_{\mathcal{N}}^{<0})$. By Proposition 2.13, we have $\mathcal{M} = \mathcal{N}$. □

To prove Propositions 2.13 and 2.14, we need the following result.

Lemma 2.15. (a) *For any subcategory $\mathcal{M}$ of $\mathcal{T}$, we have*

$$\mathsf{thick}\mathcal{M} = \bigcup_{\ell>0,\ n_1,\cdots,n_\ell \in \mathbb{Z}} \mathsf{smd}(\mathcal{M}[n_1] * \cdots * \mathcal{M}[n_\ell]).$$

(b) *Moreover if* $\mathrm{Hom}_{\mathcal{T}}(\mathcal{M}, \mathcal{M}[>0]) = 0$, *then we have*

$$\mathsf{thick}\mathcal{M} = \bigcup_{\ell \geq 0} \mathsf{smd}(\mathcal{M}[-\ell] * \mathcal{M}[1-\ell] * \cdots * \mathcal{M}[\ell-1] * \mathcal{M}[\ell]).$$

Proof. (a) Clearly the right hand side is contained in $\mathsf{thick}\mathcal{M}$. Since the right hand side clearly forms a thick subcategory of $\mathcal{T}$, it contains $\mathsf{thick}\mathcal{M}$.

(b) If $n \geq m$, then any triangle

$$M[n] \to X \to M'[m] \xrightarrow{f} M[n+1]$$

with $M, M' \in \mathcal{M}$ satisfies $f = 0$. This implies $X \simeq M[n] \oplus M'[m]$, so we have $\mathcal{M}[n] * \mathcal{M}[m] \subset \mathcal{M}[m] * \mathcal{M}[n]$. Thus we have reduced to the assertion from (a). □

The following observation is clear since we only consider subcategories $\mathcal{M}$ satisfying $\mathcal{M} = \mathsf{add}\mathcal{M}$ by our notations.

Lemma 2.16. *Let* $\mathcal{M}$ *and* $\mathcal{N}$ *be subcategories of* $\mathcal{T}$ *and* $X \in \mathsf{smd}(\mathcal{M} * \mathcal{N})$.

(a) *If* $\mathrm{Hom}_{\mathcal{T}}(\mathcal{M}, X) = 0$, *then* $X \in \mathcal{N}$.

(b) *If* $\mathrm{Hom}_{\mathcal{T}}(X, \mathcal{N}) = 0$, *then* $X \in \mathcal{M}$.

Now we have the following description of $\mathcal{T}^{\leq 0}_{\mathcal{M}}$.

Proposition 2.17. *For any* $\mathcal{M} \in \mathrm{silt}\,\mathcal{T}$, *we have*

$$\begin{aligned} \mathcal{T} &= \bigcup_{\ell \geq 0} \mathsf{smd}(\mathcal{M}[-\ell] * \mathcal{M}[1-\ell] * \cdots * \mathcal{M}[\ell-1] * \mathcal{M}[\ell]), \\ \mathcal{T}^{\leq 0}_{\mathcal{M}} &= \bigcup_{\ell \geq 0} \mathsf{smd}(\mathcal{M} * \mathcal{M}[1] * \cdots * \mathcal{M}[\ell]). \end{aligned}$$

Proof. The first equation is clear from Lemma 2.15(b).

Fix any $X \in \mathcal{T}^{\leq 0}_{\mathcal{M}}$. We can take the smallest integer n such that

$$X \in \mathsf{smd}(\mathcal{M}[n] * \mathcal{M}[n+1] \cdots * \mathcal{M}[\ell])$$

for some $\ell \geq n$. By Lemma 2.16, the minimality of n implies $\mathrm{Hom}_{\mathcal{T}}(\mathcal{M}[n], X) \neq 0$. Since $X \in \mathcal{T}^{\leq 0}_{\mathcal{M}}$, we have $n \geq 0$. Thus we have $X \in \mathsf{smd}(\mathcal{M} * \mathcal{M}[1] * \cdots * \mathcal{M}[\ell])$. □

Now we have the following important property of silting subcategories.

Theorem 2.18. *If* $\mathcal{M}, \mathcal{N} \in \mathrm{silt}\,\mathcal{T}$ *satisfy* $\mathcal{M} \subset \mathcal{N}$, *then* $\mathcal{M} = \mathcal{N}$.

Proof. Fix $X \in \mathcal{N}$. Since $\mathcal{N}$ is silting, we have $X \in \mathcal{T}^{\leq 0}_{\mathcal{M}}$. By Proposition 2.17, we have

$$X \in \mathsf{smd}(\mathcal{M} * \mathcal{M}[1] * \cdots * \mathcal{M}[\ell])$$

for some $\ell \geq 0$. Since $\mathrm{Hom}_{\mathcal{T}}(X, \mathcal{M}[1] * \cdots * \mathcal{M}[\ell]) = 0$, we have $X \in \mathcal{M}$ by Lemma 2.16. □

Now we give a proof of Proposition 2.13.

Put $\mathcal{N} := \mathcal{T}^{\leq 0}_{\mathcal{M}} \cap {}^{\perp}(\mathcal{T}^{<0}_{\mathcal{M}})$. We have $\mathcal{M} \subset \mathcal{T}^{\leq 0}_{\mathcal{M}}$. Since we have $\mathrm{Hom}_{\mathcal{T}}(\mathcal{M}, \mathcal{T}^{<0}_{\mathcal{M}}) = 0$, we have $\mathcal{M} \subset {}^{\perp}(\mathcal{T}^{<0}_{\mathcal{M}})$. Thus we have $\mathcal{M} \subset \mathcal{N}$.

On the other hand, we have

$$\mathrm{Hom}_{\mathcal{T}}(\mathcal{N}, \mathcal{N}[>0]) \subset \mathrm{Hom}_{\mathcal{T}}({}^{\perp}(\mathcal{T}^{<0}_{\mathcal{M}}), \mathcal{T}^{\leq 0}_{\mathcal{M}}[>0]) = 0.$$

Consequently, $\mathcal{N}$ is also a silting subcategory of $\mathcal{T}$. By Theorem 2.18, we have $\mathcal{M} = \mathcal{N}$. □

Now we give a proof of Proposition 2.14.

First we assume $\mathcal{T}^{\leq 0}_{\mathcal{M}} \supset \mathcal{T}^{\leq 0}_{\mathcal{N}}$. Then we have $\mathcal{N} \subset \mathcal{T}^{\leq 0}_{\mathcal{M}}$, which implies $\mathrm{Hom}_{\mathcal{T}}(\mathcal{M}, \mathcal{N}[>0]) = 0$.

Conversely we assume $\mathcal{M} \geq \mathcal{N}$. Then we have $\mathcal{N}[\geq 0] \subset \mathcal{T}^{\leq 0}_{\mathcal{M}}$. By Proposition 2.17, we have

$$\mathcal{T}^{\leq 0}_{\mathcal{N}} = \bigcup_{\ell \geq 0} \mathsf{smd}(\mathcal{N} * \mathcal{N}[1] * \cdots * \mathcal{N}[\ell]) \subset \mathcal{T}^{\leq 0}_{\mathcal{M}}.$$

Thus we have completed the proof. □

Now we give the following property of the partial order.

Proposition 2.19. *If $\mathcal{M},\mathcal{N},\mathcal{L} \in \operatorname{silt}\mathcal{T}$ satisfy $\mathcal{M} \geq \mathcal{L} \geq \mathcal{N}$, then $\mathcal{M} \cap \mathcal{N} \subset \mathcal{L}$.*

Proof. By Proposition 2.14, we have $\mathcal{T}_{\mathcal{M}}^{\leq 0} \supset \mathcal{T}_{\mathcal{L}}^{\leq 0} \supset \mathcal{T}_{\mathcal{N}}^{\leq 0}$ and ${}^{\perp}(\mathcal{T}_{\mathcal{M}}^{<0}) \subset {}^{\perp}(\mathcal{T}_{\mathcal{L}}^{<0}) \subset {}^{\perp}(\mathcal{T}_{\mathcal{N}}^{<0})$. By Proposition 2.13, we have

$$\mathcal{M} \cap \mathcal{N} \subset {}^{\perp}(\mathcal{T}_{\mathcal{M}}^{\leq 0}) \cap \mathcal{T}_{\mathcal{N}}^{\leq 0} \subset {}^{\perp}(\mathcal{T}_{\mathcal{L}}^{\leq 0}) \cap \mathcal{T}_{\mathcal{L}}^{\leq 0} = \mathcal{L}.$$

□

We end this subsection by the following observation.

Proposition 2.20. *If $\mathcal{T}$ has a silting object, then any silting subcategory contains an additive generator.*

Proof. Let M be a silting object and $\mathcal{N}$ be a silting subcategory. By Lemma 2.15, there exists $N_1,\dots,N_\ell \in \mathcal{N}$ and $n_1,\cdots,n_\ell \in \mathbb{Z}$ such that $M \in \mathsf{smd}(N_1[n_1] * \cdots * N_\ell[n_\ell])$. Let $\mathcal{N}' := \mathsf{add}(N_1 \oplus \cdots \oplus N_\ell)$. Since $M \in \mathsf{thick}\mathcal{N}'$, we have $\mathcal{T} = \mathsf{thick}\mathcal{N}'$. Thus $\mathcal{N}'$ is a silting subcategory of $\mathcal{T}$. By Theorem 2.18, we have $\mathcal{N} = \mathcal{N}'$. Thus $\mathcal{N}$ has an additive generator. □

2.3. **Krull-Schmidt triangulated categories.** Let $\mathcal{T}$ be a triangulated category. In this subsection we always assume that $\mathcal{T}$ is *Krull-Schmidt* in the sense that any object in $\mathcal{T}$ is isomorphic to a finite coproduct of objects whose endomorphism rings are local. In this case such a coproduct is uniquely determined up to isomorphism. We denote by $J_{\mathcal{T}}$ the *Jacobson radical* of $\mathcal{T}$ [ARS, ASS]. For a subcategory $\mathcal{M}$ of $\mathcal{T}$, we denote by $\mathsf{ind}\mathcal{M}$ the set of isoclasses of indecomposable objects in $\mathcal{M}$.

We say that an object $M \in \mathcal{T}$ is *basic* if M is isomorphic to a coproduct of indecomposable objects which are mutually non-isomorphic. Since $\mathcal{T}$ is Krull-Schmidt, we have a one-to-one correspondence between the isomorphism classes of basic objects M and subcategories $\mathcal{M}$ of $\mathcal{T}$ containing additive generators. It is given by $M \mapsto \mathcal{M} = \mathsf{add}M$.

Proposition 2.21. *Assume that $\mathcal{T}$ has a silting object. Then we can regard $\operatorname{silt}\mathcal{T}$ as the set of isomorphism classes of basic silting objects in $\mathcal{T}$.*

Proof. $\operatorname{silt}\mathcal{T}$ is a set by Remark 2.9. By Proposition 2.20, any silting subcategory of $\mathcal{T}$ is an additive closure of a silting object. Thus we have the assertion. □

Thanks to Krull-Schmidt assumption, we have the following useful property (e.g. [IY, 2.1, 2.3]), where (b) and (c) is famous as a 'Wakamatsu's Lemma'.

Lemma 2.22. *Let $\mathcal{M}$ be a subcategory of $\mathcal{T}$. Then the following statements hold.*

(a) *If a subcategory $\mathcal{N}$ of $\mathcal{T}$ satisfies $\operatorname{Hom}_{\mathcal{T}}(\mathcal{M},\mathcal{N}) = 0$, then $\mathcal{M} * \mathcal{N}$ is closed under summands.*
(b) *If $\mathcal{M}$ is contravariantly finite and $\mathcal{M} * \mathcal{M} \subset \mathcal{M}$, then $(\mathcal{M},\mathcal{M}^{\perp})$ is a torsion pair.*
(c) *If $\mathcal{M}$ is covariantly finite and $\mathcal{M} * \mathcal{M} \subset \mathcal{M}$, then $({}^{\perp}\mathcal{M},\mathcal{M})$ is a torsion pair.*

Now we have the following equalities.

Proposition 2.23. (a) *For any $\mathcal{M} \in \operatorname{silt}\mathcal{T}$, we have*

$$\begin{aligned}
\mathcal{T} &= \bigcup_{\ell \geq 0} \mathcal{M}[-\ell] * \mathcal{M}[1-\ell] * \cdots * \mathcal{M}[\ell-1] * \mathcal{M}[\ell], \\
\mathcal{T}_{\mathcal{M}}^{\leq 0} &= \bigcup_{\ell \geq 0} \mathcal{M} * \mathcal{M}[1] * \cdots * \mathcal{M}[\ell], \\
{}^{\perp}(\mathcal{T}_{\mathcal{M}}^{\leq 0}) &= \bigcup_{\ell > 0} \mathcal{M}[-\ell] * \mathcal{M}[-\ell+1] * \cdots * \mathcal{M}[-1].
\end{aligned}$$

(b) $({}^{\perp}(\mathcal{T}_{\mathcal{M}}^{\leq 0}), \mathcal{T}_{\mathcal{M}}^{\leq 0})$ *is a torsion pair, and so $({}^{\perp}(\mathcal{T}_{\mathcal{M}}^{<0}), \mathcal{T}_{\mathcal{M}}^{\leq 0})$ is a co-t-structure with the coheart $\mathcal{M}$.*

Proof. The first two equalities follow from Proposition 2.17 and Lemma 2.22(a). By the second equality we have

$$^{\perp}(\mathcal{T}_{\mathcal{M}}^{\leq 0}) \supset \bigcup_{\ell>0} \mathcal{M}[-\ell] * \mathcal{M}[-\ell+1] * \cdots * \mathcal{M}[-1].$$

Together with the first equality we have

$$(\bigcup_{\ell>0} \mathcal{M}[-\ell] * \cdots * \mathcal{M}[-1]) * \mathcal{T}_{\mathcal{M}}^{\leq 0} = (\bigcup_{\ell>0} \mathcal{M}[-\ell] * \cdots * \mathcal{M}[-1]) * (\bigcup_{\ell>0} \mathcal{M} * \cdots \mathcal{M}[\ell]) = \mathcal{T}.$$

Thus $(\bigcup_{\ell>0} \mathcal{M}[-\ell] * \cdots * \mathcal{M}[-1], \mathcal{T}_{\mathcal{M}}^{\leq 0})$ is a torsion pair, and hence we have

$$^{\perp}(\mathcal{T}_{\mathcal{M}}^{\leq 0}) \subset \bigcup_{\ell>0} \mathcal{M}[-\ell] * \mathcal{M}[-\ell+1] * \cdots * \mathcal{M}[-1].$$

Thus we get the third equality, and $(^{\perp}(\mathcal{T}_{\mathcal{M}}^{\leq 0}), \mathcal{T}_{\mathcal{M}}^{\leq 0})$ is a torsion pair. The coheart of the co-t-structure $(^{\perp}(\mathcal{T}_{\mathcal{M}}^{<0}), \mathcal{T}_{\mathcal{M}}^{\leq 0})$ is $\mathcal{M}$ by Proposition 2.13. □

Proposition 2.24. *Let $\mathcal{M} \in \operatorname{silt} \mathcal{T}$. For any $N = N_0 \in \mathcal{T}_{\mathcal{M}}^{\leq 0}$, we have triangles*

$$\begin{aligned}
&N_1 \xrightarrow{g_1} M_0 \xrightarrow{f_0} N_0 \longrightarrow N_1[1],\\
&\cdots,\\
&N_\ell \xrightarrow{g_\ell} M_{\ell-1} \xrightarrow{f_{\ell-1}} N_{\ell-1} \longrightarrow N_\ell[1],\\
&0 \xrightarrow{g_{\ell+1}} M_\ell \xrightarrow{f_\ell} N_\ell \longrightarrow 0,
\end{aligned}$$

for some $\ell \geq 0$ such that f_i is a minimal right $\mathcal{M}$-approximation and g_{i+1} belongs to $J_{\mathcal{T}}$ for any $0 \leq i \leq \ell$.

Proof. Since $N_0 \in \mathcal{T}_{\mathcal{M}}^{\leq 0}$, we have

$$N_0 \in \mathcal{M} * \mathcal{M}[1] * \cdots * \mathcal{M}[\ell]$$

for some $\ell \geq 0$ by Proposition 2.23. We can assume $\ell > 0$. Then we have a triangle

$$N_1' \to M_0' \xrightarrow{f_0'} N_0 \to N_1'[1]$$

with $M_0' \in \mathcal{M}$ and $N_1' \in \mathcal{M} * \mathcal{M}[1] * \cdots * \mathcal{M}[\ell-1]$. Since $N_1' \in \mathcal{T}_{\mathcal{M}}^{\leq 0}$, we have that f_0' is a right $\mathcal{M}$-approximation. Thus we can write $f_0' = (f_0\ 0) : M_0 = M_0' \oplus M_0'' \to N_0$ with a minimal right $\mathcal{M}$-approximation f_0. Then we have a triangle

$$N_1 \xrightarrow{g_1} M_0 \xrightarrow{f_0} N_0 \to N_1[1]$$

such that g_1 belongs to $J_{\mathcal{T}}$ and N_1 is a summand of N_1'. By Lemma 2.22, we have $N_1 \in \mathcal{M} * \mathcal{M}[1] * \cdots * \mathcal{M}[\ell-1]$. Repeating similar construction, we obtain the desired triangles. □

The following observation is often useful.

Lemma 2.25. *Let $\mathcal{M} \in \operatorname{silt} \mathcal{T}$ and $N_0, N_0' \in \mathcal{T}_{\mathcal{M}}^{\leq 0}$. For N_0, we take $\ell \geq 0$ and triangles in Proposition 2.24. Also for N_0', we take triangles*

$$\begin{aligned}
&N_1' \xrightarrow{g_1'} M_0' \xrightarrow{f_0'} N_0' \longrightarrow N_1'[1],\\
&\cdots,\\
&N_{\ell'}' \xrightarrow{g_{\ell'}'} M_{\ell'-1}' \xrightarrow{f_{\ell'-1}'} N_{\ell'-1}' \longrightarrow N_{\ell'}'[1],\\
&0 \xrightarrow{g_{\ell'+1}'} M_{\ell'}' \xrightarrow{f_{\ell'}'} N_{\ell'}' \longrightarrow 0,
\end{aligned}$$

satisfying the same properties. If $\operatorname{Hom}_{\mathcal{T}}(N_0, N_0'[\ell]) = 0$ holds, then we have $(\mathsf{add} M_\ell) \cap (\mathsf{add} M_0') = 0$.

Proof. For $\ell = 0$, the assertion follows from right minimality of f'_0. So we assume $\ell > 0$. We only have to show that any morphism $a : M_\ell \to M'_0$ belongs to $J_\mathcal{T}$. Applying $\mathrm{Hom}_\mathcal{T}(-, N'_0)$ to triangles in Proposition 2.24, we have

$$\mathrm{Hom}_\mathcal{T}(N_{\ell-1}[-1], N'_0) \simeq \mathrm{Hom}_\mathcal{T}(N_{\ell-2}[-2], N'_0) \simeq \cdots \simeq \mathrm{Hom}_\mathcal{T}(N_0[-\ell], N'_0) = 0.$$

Thus we have the following commutative diagram of triangles.

$$\begin{array}{ccccccccc}
N_{\ell-1}[-1] & \longrightarrow & M_\ell & \xrightarrow{g_\ell} & M_{\ell-1} & \xrightarrow{f_{\ell-1}} & N_{\ell-1} & & \\
\downarrow & & \downarrow{\scriptstyle a} & & \downarrow{\scriptstyle b} & & \downarrow & & \\
N'_1 & \xrightarrow{g'_1} & M'_0 & \xrightarrow{f'_0} & N'_0 & \longrightarrow & N'_1[1]. & &
\end{array}$$

Since f'_0 is a right $\mathcal{M}$-approximation, there exists $b' : M_{\ell-1} \to M'_0$ such that $b = f'_0 b'$. Since $f'_0(a - b'g_\ell) = 0$, there exists $a' : M_\ell \to N'_1$ such that $a = b'g_\ell + g'_1 a'$. Since both g_ℓ and g'_1 belong to $J_\mathcal{T}$, we have $a \in J_\mathcal{T}$. Thus the proof is completed. □

As an application, we have the following result.

Theorem 2.26. *If $\mathcal{T}$ has an indecomposable silting object M, then we have* $\mathrm{silt}\,\mathcal{T} = \{M[i] \mid i \in \mathbb{Z}\}$.

Proof. Let N be a basic silting object in $\mathcal{T}$. Take the smallest integer $k \in \mathbb{Z}$ such that $\mathrm{Hom}_\mathcal{T}(M, N[k]) \neq 0$. Replacing N by $N[k]$, we can assume $N \in \mathcal{T}_M^{\leq 0}$ and $\mathrm{Hom}_\mathcal{T}(M, N) \neq 0$. We have triangles in Proposition 2.24. Then we have $M_0 \neq 0$ since $\mathrm{Hom}_\mathcal{T}(M, N) \neq 0$, and moreover we can assume $M_\ell \neq 0$.

Assume $\ell > 0$. Then we have $\mathrm{Hom}_\mathcal{T}(N, N[\ell]) = 0$ and so $\mathsf{add}M_\ell \cap \mathsf{add}M_0 = 0$ by Lemma 2.25. Since both M_0 and M_ℓ are non-zero objects in $\mathsf{add}M$ where M is indecomposable, this is a contradiction. Thus we have $\ell = 0$ and $N \simeq M_0 \in \mathsf{add}M$. Since N is basic, we have $N \simeq M$. □

Next we shall show the following description of Grothendieck groups of triangulated categories with silting subcategories.

Theorem 2.27. *Let $\mathcal{T}$ be a Krull-Schmidt triangulated category with a silting subcategory $\mathcal{M}$. Then the Grothendieck group $K_0(\mathcal{T})$ of $\mathcal{T}$ is a free abelian group with a basis* $\mathsf{ind}\mathcal{M}$.

For an object $X \in \mathcal{T}$, we denote by $\delta(X)$ the number of non-isomorphic indecomposable summands of X. As an immediate consequence of Theorem 2.27, we have the following result.

Corollary 2.28. *For any silting objects $M, N \in \mathcal{T}$, we have $\delta(M) = \delta(N)$.*

Let us start with proving Theorem 2.27.

Since $\mathsf{thick}\mathcal{M} = \mathcal{T}$, we have that $\mathsf{ind}\mathcal{M}$ generate $K_0(\mathcal{T})$. It is enough to show that they are linearly independent. To prove this, we shall define a map

$$\gamma : \mathrm{ob}\mathcal{T} \to \mathbb{Z}^{\mathsf{ind}\mathcal{M}}.$$

- The map γ naturally identifies $\mathrm{ob}\mathcal{M}$ with $\mathbb{Z}^{\mathsf{ind}\mathcal{M}}_{\geq 0}$.
- For any $N \in \mathcal{T}_\mathcal{M}^{\leq 0}$, we take triangles in Proposition 2.24 and put

$$\gamma(N) := \sum_{i=0}^{\ell} (-1)^i \gamma(M_i).$$

- For general $N \in \mathcal{T}$, take a sufficiently large k such that $N[k] \in \mathcal{T}_\mathcal{M}^{\leq 0}$ and put

$$\gamma(N) := (-1)^k \gamma(N[k]).$$

In other words, for any $\ell \geq 0$ and $M_i \in \mathcal{M}$, we put

$$\gamma(M_{-\ell}[-\ell] * M_{1-\ell}[-\ell] * \cdots * M_{\ell-1}[\ell - 1] * M_\ell[\ell]) := \sum_{i=-\ell}^{\ell} (-1)^i \gamma(M_i).$$

The crucial step is to prove the following observation.

Lemma 2.29. *Let $X \to Y \to Z \to X[1]$ be a triangle in $\mathcal{T}$. Then we have $\gamma(X) - \gamma(Y) + \gamma(Z) = 0$.*

Proof. Without loss of generality, we can assume $X, Y, Z \in \mathcal{T}_{\mathcal{M}}^{\leq 0}$. To use induction, we consider the following assertions for $\ell \geq 0$.

(i)$_\ell$ The statement is true if $X, Y, Z \in \mathcal{M} * \mathcal{M}[1] * \cdots * \mathcal{M}[\ell]$.

(ii)$_\ell$ The statement is true if $X, Y \in \mathcal{M} * \mathcal{M}[1] * \cdots * \mathcal{M}[\ell]$ and $Z \in \mathcal{M} * \mathcal{M}[1] * \cdots * \mathcal{M}[\ell+1]$.

(iii)$_\ell$ The statement is true if $X \in \mathcal{M} * \mathcal{M}[1] * \cdots * \mathcal{M}[\ell]$ and $Y, Z \in \mathcal{M} * \mathcal{M}[1] * \cdots * \mathcal{M}[\ell+1]$.

The statement (i)$_0$ is true since any triangle $X \to Y \to Z \to X[1]$ with $X, Y, Z \in \mathcal{M}$ splits, so we have $Y \simeq X \coprod Z$ and $\gamma(X) - \gamma(Y) + \gamma(Z) = 0$.

We shall show (i)$_\ell \Rightarrow$(ii)$_\ell$ for any $\ell \geq 0$.

Let $X \to Y \to Z \to X[1]$ be a triangle with $X, Y \in \mathcal{M} * \mathcal{M}[1] * \cdots * \mathcal{M}[\ell]$ and $Z \in \mathcal{M} * \mathcal{M}[1] * \cdots * \mathcal{M}[\ell+1]$. Take a triangle $Z_1 \to M_0 \to Z \to Z_1[1]$ such that $M_0 \in \mathcal{M}$, $Z_1 \in \mathcal{M} * \mathcal{M}[1] * \cdots \mathcal{M}[\ell]$ and

$$\gamma(Z) = \gamma(M_0) - \gamma(Z_1). \tag{1}$$

By octahedral axiom, we have the following commutative diagram of triangles:

$$\begin{array}{ccccccc}
 & & Z_1[1] & = & Z_1[1] & & \\
 & & \uparrow & & \uparrow & & \\
X & \longrightarrow & Y & \longrightarrow & Z & \longrightarrow & X[1] \\
\| & & \uparrow & & \uparrow & & \| \\
X & \longrightarrow & W & \longrightarrow & M_0 & \longrightarrow & X[1] \\
 & & \uparrow & & \uparrow & & \\
 & & Z_1 & = & Z_1 & &
\end{array}$$

Since $\mathrm{Hom}_{\mathcal{T}}(M_0, X[1]) = 0$, the lower horizontal triangle splits and we have $W \simeq X \oplus M_0$. In particular, we have

$$\gamma(W) = \gamma(X) + \gamma(M_0). \tag{2}$$

Since the left vertical triangle $Z_1 \to W \to Y \to Z_1[1]$ satisfies $Z_1, W, Y \in \mathcal{M} * \mathcal{M}[1] * \cdots * \mathcal{M}[\ell]$, our assumption (i)$_\ell$ implies

$$\gamma(Z_1) - \gamma(W) + \gamma(Y) = 0. \tag{3}$$

Using (1), (2) and (3), we have

$$\gamma(X) - \gamma(Y) + \gamma(Z) = (\gamma(Z) - \gamma(M_0) + \gamma(Z_1)) + (\gamma(X) - \gamma(W) + \gamma(M_0)) - (\gamma(Z_1) - \gamma(W) + \gamma(Y)) = 0.$$

Thus (ii)$_\ell$ holds.

By a quite similar argument, one can show (ii)$_\ell \Rightarrow$(iii)$_\ell \Rightarrow$(i)$_{\ell+1}$ for any $\ell \geq 0$. Thus the assertion follows inductively. □

Now we are ready to prove Theorem 2.27.

By Lemma 2.29, we have a homomorphism $\gamma : K_0(\mathcal{T}) \to \mathbb{Z}^{\mathsf{ind}\mathcal{M}}$. Since $\gamma(\mathsf{ind}\mathcal{M})$ is a basis of $\mathbb{Z}^n$, the set $\mathsf{ind}\mathcal{M}$ must be linearly independent in $K_0(\mathcal{T})$. Thus it forms a basis of $K_0(\mathcal{T})$. □

2.4. **Silting mutation.** The aim of this subsection is to introduce silting mutation and give its basic properties. Let $\mathcal{T}$ be a triangulated category. We do not assume anything else on $\mathcal{T}$ unless otherwise stated.

Definition 2.30. Let $\mathcal{M} \in \mathrm{silt}\,\mathcal{T}$. For a covariantly finite subcategory $\mathcal{D}$ of $\mathcal{M}$, we define a subcategory $\mu^+(\mathcal{M}; \mathcal{D})$ of $\mathcal{T}$ as follows: For any $M \in \mathcal{M}$ we take a left $\mathcal{D}$-approximation $f : M \to D$ and a triangle

$$M \xrightarrow{f} D \xrightarrow{g} N_M \to M[1]. \tag{4}$$

We put

$$\mu^+(\mathcal{M}; \mathcal{D}) \quad := \quad \mathsf{add}(\mathcal{D} \cup \{N_M \mid M \in \mathcal{M}\}).$$

It is easily checked that $\mu^+(\mathcal{M}; \mathcal{D})$ does not depend on a choice of left approximation f. We call $\mu^+(\mathcal{M}; \mathcal{D})$ a *left mutation* of $\mathcal{M}$. Dually, we define a *right mutation* $\mu^-(\mathcal{M}; \mathcal{D})$ for a contravariantly finite subcategory $\mathcal{D}$ of $\mathcal{M}$. *(Silting) mutation* is a left or right mutation.

Theorem 2.31. *Any mutation of a silting subcategory is again a silting subcategory.*

Proof. The triangle (4) shows $\mathsf{thick}\mu^+(\mathcal{M};\mathcal{D}) \supset \mathsf{thick}\mathcal{M} = \mathcal{T}$.

We only have to show $\mathrm{Hom}_{\mathcal{T}}(\mu^+(\mathcal{M};\mathcal{D}), \mu^+(\mathcal{M};\mathcal{D})[>0]) = 0$.

Applying $\mathrm{Hom}_{\mathcal{T}}(\mathcal{D}, -[>0])$ to (4), we have an exact sequence

$$0 = \mathrm{Hom}_{\mathcal{T}}(\mathcal{D}, D[>0]) \to \mathrm{Hom}_{\mathcal{T}}(\mathcal{D}, N_M[>0]) \to \mathrm{Hom}_{\mathcal{T}}(\mathcal{D}, M[>1]) = 0$$

and so $\mathrm{Hom}_{\mathcal{T}}(\mathcal{D}, N_M[>0]) = 0$. Applying $\mathrm{Hom}_{\mathcal{T}}(-, \mathcal{D}[>0])$ to (4), we have an exact sequence

$$\mathrm{Hom}_{\mathcal{T}}(D[1], \mathcal{D}[>0]) \xrightarrow{\cdot f[1]} \mathrm{Hom}_{\mathcal{T}}(M[1], \mathcal{D}[>0]) \to \mathrm{Hom}_{\mathcal{T}}(N_M, \mathcal{D}[>0]) \to \mathrm{Hom}_{\mathcal{T}}(D, \mathcal{D}[>0]) = 0.$$

Since the above $(\cdot f[1])$ is surjective, we have $\mathrm{Hom}_{\mathcal{T}}(N_M, \mathcal{D}[>0]) = 0$. Applying $\mathrm{Hom}_{\mathcal{T}}(-, M[>1])$ to (4), we have an exact sequence

$$0 = \mathrm{Hom}_{\mathcal{T}}(M[1], M[>1]) \to \mathrm{Hom}_{\mathcal{T}}(N_M, M[>1]) \to \mathrm{Hom}_{\mathcal{T}}(D, M[>1]) = 0$$

and so $\mathrm{Hom}_{\mathcal{T}}(N_M, M[>1]) = 0$. Applying $\mathrm{Hom}_{\mathcal{T}}(N_M, -[>0])$ to (4), we have an exact sequence

$$0 = \mathrm{Hom}_{\mathcal{T}}(N_M, D[>0]) \to \mathrm{Hom}_{\mathcal{T}}(N_M, N_M[>0]) \to \mathrm{Hom}_{\mathcal{T}}(N_M, M[>1]) = 0$$

and so we have $\mathrm{Hom}_{\mathcal{T}}(N_M, N_M[>0]) = 0$.

Consequently we have $\mathrm{Hom}_{\mathcal{T}}(\mu^+(\mathcal{M};\mathcal{D}), \mu^+(\mathcal{M};\mathcal{D})[>0]) = 0$. □

In general silting mutation of a tilting subcategory is not necessarily a tilting subcategory. For this, we have the following criterion.

Theorem 2.32. *Let $\mathcal{M}$ be a tilting subcategory of $\mathcal{T}$.*

(a) *For a covariantly finite subcategory $\mathcal{D}$ of $\mathcal{M}$, the following conditions are equivalent.*
   (i) *$\mu^+(\mathcal{M};\mathcal{D})$ is tilting.*
   (ii) *Any $M \in \mathcal{M}$ has a left $\mathcal{D}$-approximation f such that $\mathrm{Hom}_{\mathcal{T}}(\mathcal{D}, f)$ is injective.*

(b) *For a contravariantly finite subcategory $\mathcal{D}$ of $\mathcal{M}$, the following conditions are equivalent.*
   (i) *$\mu^-(\mathcal{M};\mathcal{D})$ is tilting.*
   (ii) *Any $M \in \mathcal{M}$ has a right $\mathcal{D}$-approximation g such that $\mathrm{Hom}_{\mathcal{T}}(g, \mathcal{D})$ is injective.*

In these cases mutation is called tilting mutation.

Proof. We only prove the statement (a). The proof is parallel to that of Theorem 2.31.

Applying $\mathrm{Hom}_{\mathcal{T}}(\mathcal{D}, -[\neq 0])$ to (4), we have an exact sequence

$$0 = \mathrm{Hom}_{\mathcal{T}}(\mathcal{D}, D[\neq 0]) \to \mathrm{Hom}_{\mathcal{T}}(\mathcal{D}, N_M[\neq 0]) \to \mathrm{Hom}_{\mathcal{T}}(\mathcal{D}, M[\neq 1]) \xrightarrow{f\cdot} \mathrm{Hom}_{\mathcal{T}}(\mathcal{D}, D[\neq 1]).$$

Thus $\mathrm{Hom}_{\mathcal{T}}(\mathcal{D}, N_M[\neq 0]) = 0$ holds if and only if (ii) holds.

Applying $\mathrm{Hom}_{\mathcal{T}}(-, \mathcal{D}[\neq 0])$ to (4), we have an exact sequence

$$\mathrm{Hom}_{\mathcal{T}}(D[1], \mathcal{D}[\neq 0]) \xrightarrow{\cdot f[1]} \mathrm{Hom}_{\mathcal{T}}(M[1], \mathcal{D}[\neq 0]) \to \mathrm{Hom}_{\mathcal{T}}(N_M, \mathcal{D}[\neq 0]) \to \mathrm{Hom}_{\mathcal{T}}(D, \mathcal{D}[\neq 0]) = 0.$$

Since the above $(\cdot f[1])$ is surjective, we have $\mathrm{Hom}_{\mathcal{T}}(N_M, \mathcal{D}[\neq 0]) = 0$.

Applying $\mathrm{Hom}_{\mathcal{T}}(M, -[\neq -1, 0])$ to (4), we have an exact sequence

$$0 = \mathrm{Hom}_{\mathcal{T}}(M, D[\neq -1, 0]) \to \mathrm{Hom}_{\mathcal{T}}(M, N_M[\neq -1, 0]) \to \mathrm{Hom}_{\mathcal{T}}(M, M[\neq 0, 1]) = 0$$

and so $\mathrm{Hom}_{\mathcal{T}}(M, N_M[\neq -1, 0]) = 0$. Applying $\mathrm{Hom}_{\mathcal{T}}(-, N_M[\neq 0, 1])$ to (4), we have an exact sequence

$$0 = \mathrm{Hom}_{\mathcal{T}}(M, N_M[\neq -1, 0]) \to \mathrm{Hom}_{\mathcal{T}}(N_M, N_M[\neq 0, 1]) \to \mathrm{Hom}_{\mathcal{T}}(D, N_M[\neq 0, 1]) = 0$$

and so $\mathrm{Hom}_{\mathcal{T}}(N_M, N_M[\neq 0, 1]) = 0$. Thus we have $\mathrm{Hom}_{\mathcal{T}}(N_M, N_M[\neq 0]) = 0$ by Theorem 2.31.

Consequently we have $\mathrm{Hom}_{\mathcal{T}}(\mu^+(\mathcal{M};\mathcal{D}), \mu^+(\mathcal{M};\mathcal{D})[\neq 0]) = 0$ if and only if (ii) holds. □

The following properties can be checked easily.

Proposition 2.33. *Let $\mathcal{M} \in \mathrm{silt}\,\mathcal{T}$.*

(a) *For a covariantly finite subcategory $\mathcal{D}$ of $\mathcal{M}$, we have $\mu^-(\mu^+(\mathcal{M};\mathcal{D});\mathcal{D}) = \mathcal{M}$ and $\mathcal{M} \geq \mu^+(\mathcal{M};\mathcal{D})$, where the equality holds if and only if $\mathcal{D} = \mathcal{M}$.*

(b) *For a contravariantly finite subcategory $\mathcal{D}$ of $\mathcal{M}$, we have $\mu^+(\mu^-(\mathcal{M};\mathcal{D});\mathcal{D}) = \mathcal{M}$ and $\mu^-(\mathcal{M};\mathcal{D}) \geq \mathcal{M}$, where the equality holds if and only if $\mathcal{D} = \mathcal{M}$.*

Proof. (a) Applying $\mathrm{Hom}_{\mathcal{T}}(\mathcal{D}, -)$ to (4), we have an exact sequence

$$\mathrm{Hom}_{\mathcal{T}}(\mathcal{D}, D) \xrightarrow{g_*} \mathrm{Hom}_{\mathcal{T}}(\mathcal{D}, N_M) \to \mathrm{Hom}_{\mathcal{T}}(\mathcal{D}, M[1]) = 0.$$

Thus g is a right $\mathcal{D}$-approximation. Thus we have $\mu^-(\mu^+(\mathcal{M};\mathcal{D});\mathcal{D}) = \mathcal{M}$.

Applying $\mathrm{Hom}_{\mathcal{T}}(\mathcal{M}, -[>0])$ to (4), we have an exact sequence

$$0 = \mathrm{Hom}_{\mathcal{T}}(\mathcal{M}, D[>0]) \to \mathrm{Hom}_{\mathcal{T}}(\mathcal{M}, N_M[>0]) \to \mathrm{Hom}_{\mathcal{T}}(\mathcal{M}, M[>1]) = 0$$

and so $\mathrm{Hom}_{\mathcal{T}}(\mathcal{M}, N_M[>0]) = 0$. Thus $\mathrm{Hom}_{\mathcal{T}}(\mathcal{M}, \mu^+(\mathcal{M};\mathcal{D})[>0]) = 0$ holds and we have $\mathcal{M} \geq \mu^+(\mathcal{M};\mathcal{D})$.

If $\mathcal{D} \neq \mathcal{M}$, then take $M \in \mathcal{M}\backslash\mathcal{D}$. Since f is not a split monomorphism, we have $\mathrm{Hom}_{\mathcal{T}}(N_M, M[1]) \neq 0$. Thus $N_M \notin \mathcal{M}$ holds, so we have $\mathcal{M} \neq \mu^+(\mathcal{M};\mathcal{D})$. □

In the rest of this subsection, we assume that $\mathcal{T}$ is a Krull-Schmidt triangulated category. The following notation will be often used.

Definition 2.34. Let $\mathcal{M} \in \mathrm{silt}\,\mathcal{T}$ and $\mathcal{X}$ a subcategory of $\mathcal{M}$. Define a subcategory $\mathcal{M}_{\mathcal{X}}$ of $\mathcal{M}$ by $\mathsf{ind}\mathcal{M}_{\mathcal{X}} = \mathsf{ind}\mathcal{M}\backslash\mathsf{ind}\mathcal{X}$. We put

$$\mu^{\pm}_{\mathcal{X}}(\mathcal{M}) := \mu^{\pm}(\mathcal{M};\mathcal{M}_{\mathcal{X}})$$

We say that mutation is *irreducible* if $\#\mathsf{ind}\mathcal{X} = 1$. If $\mathcal{X} = \mathsf{add}X$, we denote $\mathcal{M}_{\mathcal{X}}$ and $\mu^{\pm}_{\mathcal{X}}$ by $\mathcal{M}_X$ and $\mu^{\pm}_X$ respectively.

Now we assume the following condition:

(F) $\mathcal{T}$ is Krull-Schmidt, and for any $\mathcal{M} \in \mathrm{silt}\,\mathcal{T}$ and $X \in \mathsf{ind}\mathcal{M}$, the subcategory $\mathcal{M}_X$ is functorially finite in $\mathcal{M}$.

This is satisfied if $\mathcal{T}$ is k-linear Hom-finite and has a silting object (Proposition 2.20).

Under this condition, we shall show the following result.

Theorem 2.35. *Let $\mathcal{T}$ be a triangulated category satisfying the condition (F). For any $\mathcal{M},\mathcal{N} \in \mathrm{silt}\,\mathcal{T}$, the following conditions are equivalent.*

(a) *$\mathcal{N}$ is an irreducible left mutation of $\mathcal{M}$.*
(b) *$\mathcal{M}$ is an irreducible right mutation of $\mathcal{N}$.*
(c) *$\mathcal{M} > \mathcal{N}$ and there is no $\mathcal{L} \in \mathrm{silt}\,\mathcal{T}$ satisfying $\mathcal{M} > \mathcal{L} > \mathcal{N}$.*

The following property is crucial in our consideration.

Proposition 2.36. *If $\mathcal{M},\mathcal{N} \in \mathrm{silt}\,\mathcal{T}$ satisfy $\mathcal{M} > \mathcal{N}$, then there exists an irreducible left mutation $\mathcal{L}$ of $\mathcal{M}$ such that $\mathcal{M} > \mathcal{L} \geq \mathcal{N}$.*

Proof. We take $N_0 \in \mathcal{N}$ which does not belong to $\mathcal{M}$, then consider $\ell \geq 0$ and triangles in Proposition 2.24. Then we have $\ell > 0$. Take an indecomposable summand X of M_ℓ, and let $\mathcal{L} := \mu^+_X(\mathcal{M})$ be an irreducible left mutation. Then we have $\mathcal{L} = \mathsf{add}(\mathcal{M}_X \cup \{Y\})$, where Y is given by the triangle

$$X \xrightarrow{f} D \xrightarrow{g} Y \to X[1]$$

with a left $\mathcal{M}_X$-approximation f of X.

We only have to prove $\mathcal{L} \geq \mathcal{N}$. We need to show $\mathrm{Hom}_{\mathcal{T}}(Y,\mathcal{N}[>0]) = 0$. Since we have an exact sequence

$$0 = \mathrm{Hom}_{\mathcal{T}}(X,\mathcal{N}[>0]) \to \mathrm{Hom}_{\mathcal{T}}(Y,\mathcal{N}[>1]) \to \mathrm{Hom}_{\mathcal{T}}(D,\mathcal{N}[>1]) = 0,$$

we have $\mathrm{Hom}_{\mathcal{T}}(Y,\mathcal{N}[>1]) = 0$. Thus it remains to show $\mathrm{Hom}_{\mathcal{T}}(Y,\mathcal{N}[1]) = 0$. Since we have an exact sequence

$$\mathrm{Hom}_{\mathcal{T}}(D,\mathcal{N}) \xrightarrow{\cdot f} \mathrm{Hom}_{\mathcal{T}}(X,\mathcal{N}) \to \mathrm{Hom}_{\mathcal{T}}(Y,\mathcal{N}[1]) \to \mathrm{Hom}_{\mathcal{T}}(D,\mathcal{N}[1]) = 0,$$

we only have to show that $\mathrm{Hom}_{\mathcal{T}}(D,\mathcal{N}) \xrightarrow{\cdot f} \mathrm{Hom}_{\mathcal{T}}(X,\mathcal{N})$ is surjective.

Fix any $N_0' \in \mathcal{N}$ and $a : X \to N_0'$. We consider the diagram

$$\begin{array}{ccccccc} & & Y[-1] & \longrightarrow & X & \xrightarrow{f} & D \xrightarrow{g} Y \\ & & & & \downarrow a & & \\ N_1' & \xrightarrow{g_0'} & M_0' & \xrightarrow{f_0'} & N_0' & \longrightarrow & N_1'[1]. \end{array}$$

where the lower triangle is given in Lemma 2.25. Since f_0' is a right $\mathcal{M}$-approximation, there exists $b : X \to M_0'$ such that $a = f_0'b$. Since $(\mathsf{add}M_\ell) \cap (\mathsf{add}M_0') = 0$ holds by Lemma 2.25, we have $X \notin \mathsf{add}M_0'$ and so $M_0' \in \mathcal{M}_X$. Since f is a left $\mathcal{M}_X$-approximation, there exists $c : D \to M_0'$ such that $b = cf$. Thus we have $a = (f_0'c)f$, and we have the assertion. □

Now we are ready to prove Theorem 2.35.

(i) We shall show that (a) and (b) are equivalent.

We only have to show that (a) implies (b). Assume that $\mathcal{N}$ is an irreducible left mutation $\mu_X^+(\mathcal{M})$ of $\mathcal{M}$ with respect to $X \in \mathsf{ind}\mathcal{M}$. Take a triangle

$$X \xrightarrow{f} D \xrightarrow{g} Y \to X[1]$$

with a minimal left $\mathcal{M}_X$-approximation f. By Proposition 2.33, we have $\mathcal{M} = \mu_Y^-(\mathcal{N})$. Since g is a right $\mathcal{M}_X$-approximation, it is easy to check that Y is indecomposable. Thus $\mathcal{M}$ is an irreducible right mutation of $\mathcal{N}$.

(ii) We shall show that (a) and (c) are equivalent.

Assume that (c) is satisfied. Since $\mathcal{M} > \mathcal{N}$, there exists an irreducible left mutation $\mathcal{L}$ of $\mathcal{M}$ such that $\mathcal{M} > \mathcal{L} \geq \mathcal{N}$ by Proposition 2.36. By the condition (c), we have $\mathcal{L} = \mathcal{N}$, and we have the assertion.

Assume that (a) is satisfied. Thus $\mathcal{N} = \mu_X^+(\mathcal{M})$ for indecomposable X. Assume that there exists $\mathcal{L}$ such that $\mathcal{M} > \mathcal{L} > \mathcal{N}$. Again by Proposition 2.36, there exists a left mutation $\mathcal{K}$ of $\mathcal{M}$ such that $\mathcal{M} > \mathcal{K} \geq \mathcal{L}$. By Proposition 2.19, we have $\mathcal{M}_X = \mathcal{M} \cap \mathcal{N} \subset \mathcal{K}$. Since both of $\mathcal{N}$ and $\mathcal{K}$ are left mutation of $\mathcal{M}$ with respect to X, we have $\mathcal{N} = \mathcal{K}$, a contradiction. Thus (c) holds. □

2.5. **Silting reduction.** In this subsection, we give a reduction theorem of silting subcategories, which is an analogue of 2-Calabi-Yau reduction in cluster tilting theory [IY, 4.9]. The following result gives a bijection between silting subcategories of $\mathcal{T}$ containing a functorially finite thick subcategory $\mathcal{S}$ and silting subcategories of the quotient triangulated category $\mathcal{T}/\mathcal{S}$.

Theorem 2.37. *Let $\mathcal{T}$ be a Krull-Schmidt triangulated category, $\mathcal{S}$ a thick subcategory of $\mathcal{T}$ and $\mathcal{U} := \mathcal{T}/\mathcal{S}$. Let $F : \mathcal{T} \to \mathcal{U}$ be the canonical functor.*

(a) *If $\mathcal{S}$ is a contravariantly finite subcategory of $\mathcal{T}$, then for any $\mathcal{D} \in \operatorname{silt}\mathcal{S}$ we have an injective map*

$$\{\mathcal{M} \in \operatorname{silt}\mathcal{T} \mid \mathcal{D} \subset \mathcal{M}\} \to \operatorname{silt}\mathcal{U}, \quad \mathcal{M} \mapsto F\mathcal{M}.$$

(b) *If $\mathcal{S}$ is a functorially finite subcategory of $\mathcal{T}$, then the map in (a) is bijective.*

The first step of the proof is to consider the subcategory $\mathcal{S}^{\perp_\tau}$ of $\mathcal{T}$. Since $\mathcal{S}$ is contravariantly finite, we have a stable t-structure $(\mathcal{S}, \mathcal{S}^{\perp_\tau})$ of $\mathcal{T}$ by Lemma 2.22. Moreover, we can naturally identify $\mathcal{U}$ with the subcategory $\mathcal{S}^{\perp_\tau}$ of $\mathcal{T}$ [M].

Next we shall show the following observation.

Lemma 2.38. *Let $\mathcal{M}$ be as in Theorem 2.37(a). For any $M \in \mathcal{M}$, take a triangle*

$$S \xrightarrow{a} M \xrightarrow{b} FM \to S[1] \tag{5}$$

with $S \in \mathcal{S}$ and $FM \in \mathcal{U}$. Then $S \in \mathcal{S}_{\mathcal{D}}^{\leq 0}$ holds, where

$$({}^{\perp_{\mathcal{S}}}(\mathcal{S}_{\mathcal{D}}^{\leq 0}), \mathcal{S}_{\mathcal{D}}^{\leq 0}) = (\bigcup_{\ell>0} \mathcal{D}[-\ell] * \cdots * \mathcal{D}[-1], \bigcup_{\ell \geq 0} \mathcal{D} * \cdots * \mathcal{D}[\ell])$$

is the torsion pair in $\mathcal{S} = \mathsf{thick}\mathcal{D}$ given in Proposition 2.23.

Proof. Take a triangle

$$S^{>0} \xrightarrow{c} S \xrightarrow{d} S^{\leq 0} \to S^{>0}[1]$$

with $S^{>0} \in {}^{\perp_{\mathcal{S}}}(\mathcal{S}_{\mathcal{D}}^{\leq 0})$ and $S^{\leq 0} \in \mathcal{S}_{\mathcal{D}}^{\leq 0}$. Since $\mathrm{Hom}_{\mathcal{T}}({}^{\perp_{\mathcal{S}}}(\mathcal{S}_{\mathcal{D}}^{\leq 0}), \mathcal{M}) = 0$, we have $ac = 0$. Thus there exists a morphism $e : S^{\leq 0} \to M$ such that $a = ed$. Since $be = 0$ by $\mathrm{Hom}_{\mathcal{D}}(\mathcal{S}, \mathcal{U}) = 0$, there exists $f : S^{\leq 0} \to S$ such that $e = af$.



Since $a(1_S - fd) = 0$, we have that $1_S - fd : S \to S$ factors through $FM[-1]$. Since $\mathrm{Hom}_{\mathcal{T}}(\mathcal{S}, \mathcal{U}) = 0$ again, we have $1_S = fd$. Thus d is a split monomorphism, and we have $S \in \mathcal{S}_{\mathcal{D}}^{\leq 0}$. □

Now we are ready to prove Theorem 2.37.

(a)(i) We shall show that $F\mathcal{M}$ is a silting subcategory of $\mathcal{U}$.

We have $\mathsf{thick} F\mathcal{M} \supset F\mathsf{thick}\mathcal{M} = F\mathcal{T} = \mathcal{U}$. Thus we only have to show $\mathrm{Hom}_{\mathcal{T}}(F\mathcal{M}, F\mathcal{M}[>0]) = 0$.

Applying $\mathrm{Hom}_{\mathcal{T}}(M, -[>0])$ to (5), we have an exact sequence

$$0 \overset{\mathcal{M} \in \mathrm{silt}\,\mathcal{T}}{=} \mathrm{Hom}_{\mathcal{T}}(M, M[>0]) \to \mathrm{Hom}_{\mathcal{T}}(M, FM[>0]) \to \mathrm{Hom}_{\mathcal{T}}(M, S[>1]) \overset{\mathrm{Hom}_{\mathcal{T}}(\mathcal{M}, \mathcal{S}_{\mathcal{D}}^{\leq 0}) = 0}{=} 0$$

and so $\mathrm{Hom}_{\mathcal{T}}(M, FM[>0]) = 0$. Applying $\mathrm{Hom}_{\mathcal{T}}(-, FM[>0])$ to (5), we have an exact sequence

$$0 \overset{\mathrm{Hom}_{\mathcal{T}}(\mathcal{S}, \mathcal{U}) = 0}{=} \mathrm{Hom}_{\mathcal{T}}(S, FM[>-1]) \to \mathrm{Hom}_{\mathcal{T}}(FM, FM[>0]) \to \mathrm{Hom}_{\mathcal{T}}(M, FM[>0]) = 0$$

and so $\mathrm{Hom}_{\mathcal{T}}(FM, FM[>0]) = 0$.

(ii) We shall show that the correspondence $\mathcal{M} \mapsto F\mathcal{M}$ is injective. By Theorem 2.11, it is enough to prove $\mathcal{M} \geq \mathcal{N}$ if $F\mathcal{M} \geq F\mathcal{N}$. Let $M \in \mathcal{M}$ and $N \in \mathcal{N}$. For any $f \in \mathrm{Hom}_{\mathcal{T}}(M, N[>0])$, we have a commutative diagram

$$\begin{array}{ccccccc} S_M & \longrightarrow & M & \longrightarrow & FM & \longrightarrow & S_M[1] \\ \downarrow & & \downarrow f & & \downarrow & & \downarrow \\ S_N[>0] & \xrightarrow[a_N[>0]]{} & N[>0] & \xrightarrow[b_N[>0]]{} & FN[>0] & \longrightarrow & S_N[>1]. \end{array}$$

Since $b_N[>0] \cdot f = 0$ by our assumption, f factors through $a_N[>0] : S_N[>0] \to N[>0]$. This implies $f = 0$ since $\mathrm{Hom}_{\mathcal{T}}(M, S_N[>0]) \subset \mathrm{Hom}_{\mathcal{T}}(\mathcal{M}, \mathcal{S}_{\mathcal{D}}^{<0}) = 0$ by Lemma 2.38. Thus we have $\mathcal{M} \geq \mathcal{N}$.

(b) We shall show that the correspondence $\mathcal{M} \mapsto F\mathcal{M}$ is surjective. Fix $\mathcal{N} \in \mathrm{silt}\,\mathcal{U}$.

Since $\mathcal{S}$ is covariantly finite in $\mathcal{T}$ by our assumption and $\mathcal{S}_{\mathcal{D}}^{<0}$ is covariantly finite in $\mathcal{S}$ by Proposition 2.23, we have that $\mathcal{S}_{\mathcal{D}}^{<0}$ is covariantly finite in $\mathcal{T}$. Thus we have a torsion pair $({}^{\perp_{\mathcal{T}}}(\mathcal{S}_{\mathcal{D}}^{<0}), \mathcal{S}_{\mathcal{D}}^{\leq 0})$ in $\mathcal{T}$ by Lemma 2.22(c). For any $N \in \mathcal{N}$, take a triangle

$$S_N \to M_N \to N \to S_N[1] \tag{6}$$

with $M_N \in {}^{\perp_{\mathcal{T}}}(\mathcal{S}_{\mathcal{D}}^{<0})$ and $S_N \in \mathcal{S}_{\mathcal{D}}^{\leq 0}$. Notice that M_N is unique up to a summand in $\mathcal{D}$ since $\mathcal{S}_{\mathcal{D}}^{\leq 0} \cap {}^{\perp_{\mathcal{T}}}(\mathcal{S}_{\mathcal{D}}^{<0}) = \mathcal{S}_{\mathcal{D}}^{\leq 0} \cap {}^{\perp_{\mathcal{S}}}(\mathcal{S}_{\mathcal{D}}^{<0}) = \mathcal{D}$ by Proposition 2.13.

We shall show that $\mathcal{M} := \mathsf{add}(\mathcal{D} \cup \{M_N \mid N \in \mathcal{N}\})$ is a silting subcategory of $\mathcal{T}$. Since $\mathcal{D}[>0] \subset \mathcal{S}_{\mathcal{D}}^{<0}$, we have $\mathrm{Hom}_{\mathcal{T}}(M_N, \mathcal{D}[>0]) = 0$. Applying $\mathrm{Hom}_{\mathcal{T}}(\mathcal{D}, -[>0])$ to (6), we have an exact sequence

$$0 \overset{\mathrm{Hom}_{\mathcal{T}}(\mathcal{D}, \mathcal{S}_{\mathcal{D}}^{<0}) = 0}{=} \mathrm{Hom}_{\mathcal{T}}(\mathcal{D}, S_N[>0]) \to \mathrm{Hom}_{\mathcal{T}}(\mathcal{D}, M_N[>0]) \to \mathrm{Hom}_{\mathcal{T}}(\mathcal{D}, N[>0]) \overset{\mathrm{Hom}_{\mathcal{T}}(\mathcal{S}, \mathcal{U}) = 0}{=} 0$$

and so $\mathrm{Hom}_{\mathcal{T}}(\mathcal{D}, M_N[>0]) = 0$.

Applying $\mathrm{Hom}_{\mathcal{T}}(-, N[>0])$ to (6), we have an exact sequence

$$0 \overset{\mathcal{N} \in \mathrm{silt}\,\mathcal{U}}{=} \mathrm{Hom}_{\mathcal{T}}(N, N[>0]) \to \mathrm{Hom}_{\mathcal{T}}(M_N, N[>0]) \to \mathrm{Hom}_{\mathcal{T}}(S_N, N[>0]) \overset{\mathrm{Hom}_{\mathcal{T}}(\mathcal{S}, \mathcal{U}) = 0}{=} 0$$

and so $\mathrm{Hom}_{\mathcal{T}}(M_N, N[>0]) = 0$. Applying $\mathrm{Hom}_{\mathcal{T}}(M_N, -[>0])$ to (6), we have an exact sequence

$$0 \overset{\mathrm{Hom}_{\mathcal{T}}({}^{\perp_{\mathcal{T}}}(\mathcal{S}_{\mathcal{D}}^{\leq 0}), \mathcal{S}_{\mathcal{D}}^{\leq 0})=0}{=} \mathrm{Hom}_{\mathcal{T}}(M_N, S_N[>0]) \to \mathrm{Hom}_{\mathcal{T}}(M_N, M_N[>0]) \to \mathrm{Hom}_{\mathcal{T}}(M_N, N[>0]) = 0$$

and so $\mathrm{Hom}_{\mathcal{T}}(M_N, M_N[>0]) = 0$.

Consequently we have $\mathrm{Hom}_{\mathcal{T}}(\mathcal{M}, \mathcal{M}[>0]) = 0$. By (6), we have $\mathsf{thick}\mathcal{M} \supset \mathcal{N}$, so we have $\mathsf{thick}\mathcal{M} \supset \mathsf{thick}(\mathcal{D} \cup \mathcal{N}) \supset \mathcal{S} * \mathcal{U} = \mathcal{T}$. Thus $\mathcal{M}$ is a silting subcategory of $\mathcal{T}$. □

The functor $F : \mathcal{M} \to F\mathcal{M}$ has the following properties.

Proposition 2.39. *In Theorem 2.37(a), the functor $F : \mathcal{M} \to F\mathcal{M}$ induces an equivalence*

$$\mathcal{M}/[\mathcal{D}] \to F\mathcal{M},$$

where $[\mathcal{D}]$ is the ideal of $\mathcal{M}$ consisting of morphisms which factor through objects in $\mathcal{D}$. In particular, F induces a bijection between $\mathsf{ind}\mathcal{M} \backslash \mathsf{ind}\mathcal{D}$ and $\mathsf{ind}F\mathcal{M}$.

Proof. Since $F\mathcal{D} = 0$, we have the induced functor $F : \mathcal{M}/[\mathcal{D}] \to F\mathcal{M}$. We only have to show that this is fully faithful. Let $M, M' \in \mathcal{M}$. For each $f : M \to M'$, the morphism $Ff : FM \to FM'$ is given by the following commutative diagram.

$$\begin{array}{ccccccc} S & \longrightarrow & M & \longrightarrow & FM & \longrightarrow & S[1] \\ \downarrow & & \downarrow{\scriptstyle f} & & \downarrow{\scriptstyle Ff} & & \downarrow \\ S' & \longrightarrow & M' & \longrightarrow & FM' & \longrightarrow & S'[1] \end{array}$$

(i) We shall show that $F : \mathcal{M}/[\mathcal{D}] \to F\mathcal{M}$ is faithful.

If $Ff = 0$, then f factors through S'. By Lemma 2.38, we know $S' \in \mathcal{S}_{\mathcal{D}}^{\leq 0} = \bigcup_{\ell \geq 0} \mathcal{D} * \cdots * \mathcal{D}[\ell]$. Since $\mathrm{Hom}_{\mathcal{T}}(M, \mathcal{D}[1] * \cdots * \mathcal{D}[\ell]) = 0$, we have that f factors through $\mathcal{D}$. Thus f is zero in $\mathcal{M}/[\mathcal{D}]$.

(ii) We shall show that $F : \mathcal{M}/[\mathcal{D}] \to F\mathcal{M}$ is full.

Fix $g \in \mathrm{Hom}_{\mathcal{T}}(FM, FM')$. Since $\mathrm{Hom}_{\mathcal{T}}(M, S'[1]) = 0$ again by Lemma 2.38, there exists $f : M \to M'$ which makes the following diagram commutative.

$$\begin{array}{ccccccc} S & \longrightarrow & M & \longrightarrow & FM & \longrightarrow & S[1] \\ \downarrow & & \downarrow{\scriptstyle f} & & \downarrow{\scriptstyle g} & & \downarrow \\ S' & \longrightarrow & M' & \longrightarrow & FM' & \longrightarrow & S'[1] \end{array}$$

Then $f = Fg$ holds. □

Later we use the following compatibility of silting mutation and silting reduction.

Lemma 2.40. *In Theorem 2.37(a), we have $F\mu_{\mathcal{X}}^{+}(\mathcal{M}) = \mu_{F\mathcal{X}}^{+}(F\mathcal{M})$ for any covariantly finite subcategory $\mathcal{X}$ of $\mathcal{M}$ satisfying $\mathcal{D} \cap \mathcal{X} = 0$.*

Proof. By Proposition 2.39, we have $(F\mathcal{M})_{F\mathcal{X}} = F(\mathcal{M}_{\mathcal{X}})$. For any $X \in \mathcal{X}$, take a triangle

$$X \xrightarrow{f} D \to Y \to X[1]$$

where f is a left $\mathcal{M}_{\mathcal{X}}$-approximation. We have to prove that Ff is a left $F(\mathcal{M}_{\mathcal{X}})$-approximation. It is enough to show $\mathrm{Hom}_{\mathcal{T}}(FY, F(\mathcal{M}_{\mathcal{X}})[1]) = 0$.

For any $M \in \mathcal{M}_{\mathcal{X}}$, there exists a triangle

$$S \to M \to FM \to S[1] \tag{7}$$

with $S \in \mathcal{S}_{\mathcal{D}}^{\leq 0}$ by Lemma 2.38. Applying $\mathrm{Hom}_{\mathcal{T}}(Y, -)$, we have an exact sequence

$$0 \overset{\text{Thm. } 2.31}{=} \mathrm{Hom}_{\mathcal{T}}(Y, M[1]) \to \mathrm{Hom}_{\mathcal{T}}(Y, FM[1]) \to \mathrm{Hom}_{\mathcal{T}}(Y, S[2]) \overset{S \in \mathcal{S}_{\mathcal{D}}^{\leq 0}}{=} 0,$$

so we have $\mathrm{Hom}_{\mathcal{T}}(Y, FM[1]) = 0$. Thus we have $\mathrm{Hom}_{\mathcal{T}}(FY, FM[1]) \simeq \mathrm{Hom}_{\mathcal{T}}(Y, FM[1]) = 0$. □

2.6. **Silting quivers and examples.** Let $\mathcal{T}$ be a Krull-Schmidt triangulated category. The aim of this section is to introduce the silting quiver of $\mathcal{T}$.

Definition 2.41. The *silting quiver* of $\mathcal{T}$ is defined as follows:

- The set of vertices is $\operatorname{silt}\mathcal{T}$.
- We draw an arrow $\mathcal{M} \to \mathcal{N}$ if $\mathcal{N}$ is an irreducible left mutation of $\mathcal{M}$.

If $\mathcal{T}$ satisfies the condition (F), then the silting quiver is nothing but the Hasse quiver of the partially ordered set $\operatorname{silt}\mathcal{T}$ by Theorem 2.35.

We pose the following question.

Question 2.42. Let A be a finite dimensional algebra A over a field and $\mathcal{T} := \mathsf{K}^{\mathrm{b}}(\text{proj-}A)$. When is the silting quiver of $\mathcal{T}$ connected? Equivalently, when is the action of iterated irreducible mutation on $\operatorname{silt}\mathcal{T}$ transitive?

If A is local, then we have a positive answer by the following observation.

Corollary 2.43. *If $\mathcal{T}$ has an indecomposable silting object, then the action of iterated irreducible mutation on* $\operatorname{silt}\mathcal{T}$ *is transitive.*

Proof. This is immediate from Theorem 2.26 since we have $\mu^{\pm}_{M[i]}(M[i]) = M[i \pm 1]$. □

Next we shall generalize Corollary 2.43 by introducing the following notion which generalizes 'almost complete partial tilting modules'.

We call a subcategory $\mathcal{D}$ of $\mathcal{T}$ *almost complete silting* if there exists a silting subcategory $\mathcal{M}$ of $\mathcal{T}$ such that $\mathcal{M} \supset \mathcal{D}$ and $\#(\mathsf{ind}\mathcal{M} \backslash \mathsf{ind}\mathcal{D}) = 1$.

Then we have the following analogue of [HU, 2.1] for 'tilting mutation' and [IY, 5.3] for 'cluster tilting mutation'.

Theorem 2.44. *Let $\mathcal{T}$ be a Krull-Schmidt triangulated category and $\mathcal{D}$ an almost complete silting subcategory. If $\mathsf{thick}\mathcal{D}$ is a functorially finite subcategory of $\mathcal{T}$, then the set $\{\mathcal{M} \in \operatorname{silt}\mathcal{T} \mid \mathcal{D} \subset \mathcal{M}\}$ is transitive under iterated irreducible mutation.*

Proof. Let $\mathcal{U} := \mathcal{T}/\mathsf{thick}\mathcal{D}$. By Theorem 2.37 we have a bijection $\{\mathcal{M} \in \operatorname{silt}\mathcal{T} \mid \mathcal{D} \subset \mathcal{M}\} \to \operatorname{silt}\mathcal{U}$. This commutes with mutation by Lemma 2.40 and its dual. By Corollary 2.43, we know that $\operatorname{silt}\mathcal{U}$ is transitive under iterated irreducible mutation. Thus the assertion follows. □

We give examples of silting quivers.

Example 2.45. Let A be a path algebra of the quiver $1 \longrightarrow 2$. We have the AR-quiver of $\mathsf{K}^{\mathrm{b}}(\text{proj-}A)$ as follows:



Then the silting quiver of A is the following (cf. Theorem 3.1):



Identifying each silting object T with $T[i]$ for any $i \in \mathbb{Z}$, we can simplify the quiver as follows, where $M \overset{[1]}{\rightarrow} N$ means $M \longrightarrow N[1]$.



Example 2.46. Let A be a path algebra of the quiver $1 \overset{\longrightarrow}{\longrightarrow} 2$ with $\ell \geq 2$ arrows. The AR-quiver of $\mathsf{K}^{\mathrm{b}}(\text{proj-}A)$ contains the following connected component:



Then the silting quiver of A is the following (cf. Theorem 3.1):



Example 2.47. Let A be an algebra presented by a quiver $1 \underset{b}{\overset{a}{\rightleftarrows}} 2$ with relations $ab = 0 = ba$. Then the silting quiver of A is the following, where $X := \mathrm{cone}(P_1 \to P_2)$ and $Y := \mathrm{cone}(P_2 \to P_1)$:



Example 2.48. Let A be an algebra presented by a quiver $1 \underset{b}{\overset{a}{\rightleftarrows}} 2$ with relations $(ab)^\ell a = 0 = (ba)^\ell b$ ($\ell \geq 1$). The silting quiver of A is connected by [A], and it is the following:



2.7. **Okuyama-Rickard complexes and APR and BB tilting modules.** Our silting mutation has two important origins in representation theory:

- Okuyama-Rickard complexes and Okuyama's method in modular representation theory;
- APR tilting modules and BB tilting modules;

In this section we will explain the relationship between silting mutation and these notions.

Throughout this section, let A be a finite dimensional algebra over a field k. We may assume that A is basic and indecomposable as an A-A-bimodule. For an A-module X, we denote by $P(X)$ a projective cover of X. We denote by $\nu = D\mathrm{Hom}_A(-, A) : \mathrm{mod}\text{-}A \to \mathrm{mod}\text{-}A$ the Nakayama functor. We denote by τ and τ^{-1} the Auslander-Reiten translations [ARS, ASS].

Definition 2.49. For idempotent $e \in A$, the *Okuyama-Rickard complex* with respect to e is defined by

$$T := \begin{cases} \overset{0}{P(eA(1-e)A)} & \xrightarrow{p_e} & \overset{1}{eA} \\ \oplus & & \\ (1-e)A & & \end{cases} \tag{8}$$

where p_e gives a projective cover of the submodule $eA(1-e)A$ of eA.

In [O], Okuyama constructed Okuyama-Rickard complexes and proved that it is tilting if A is symmetric. The method of his construction is often called *Okuyama's method*.

Let us give basic properties of Okuyama-Rickard complexes in our context of silting mutation.

Theorem 2.50. *Let $e \in A$ be an idempotent and T the Okuyama-Rickard complex with respect to e.*

(a) *T is isomorphic to right mutation $\mu_{eA}^{-}(A)$ of A with respect to eA.*
(b) *T is a silting object in $\mathsf{K}^{\mathrm{b}}(\text{proj-}A)$.*
(c) *The following conditions are equivalent.*
  (i) *T is a tilting object in $\mathsf{K}^{\mathrm{b}}(\text{proj-}A)$.*
  (ii) $\mathrm{Hom}_A(eA/eA(1-e)A, (1-e)A) = 0$.
(d) *If A is a self-injective algebra and $eA \simeq \nu(eA)$ holds, then T is a tilting object in $\mathsf{K}^{\mathrm{b}}(\text{proj-}A)$.*

Proof. (a) Since $eA(1-e)A$ is minimal amongst submodules X of eA such that any composition factor of eA/X belongs to $\mathsf{add}(\mathrm{top}\, eA)$, we have that p_e in (8) is a minimal right $\mathsf{add}(1-e)A$-approximation of eA. Thus the assertion follows.

(b)(c) These are immediate consequences of Theorems 2.31 and 2.32.

(d) This is immediate from (c) since $\mathsf{add}(\mathrm{top}\, eA) \cap \mathsf{add}(\mathrm{soc}(1-e)A) = \mathsf{add}(\mathrm{top}\, eA) \cap \mathsf{add}(\mathrm{top}(1-e)A) = 0$ holds by our assumption and any composition factor of $eA/eA(1-e)A$ is isomorphic to top eA. □

The partial order $\leq$ on $\mathrm{silt}\,\mathsf{K}^{\mathrm{b}}(\text{proj-}A)$ have the following easy interpretation for Okuyama-Rickard complexes.

Proposition 2.51. *Let $e, e' \in A$ be idempotents and T, T' the Okuyama-Rickard complexes with respect to e, e' respectively. Then the following conditions are equivalent.*

(a) $\mathsf{add}eA \supseteq \mathsf{add}e'A$.
(b) $T \geq T'$.

Proof. Assume $\mathsf{add}eA \supseteq \mathsf{add}e'A$. Clearly we have $\mathrm{Hom}_{\mathsf{K}^{\mathrm{b}}(\text{proj-}A)}(T, T'[>1]) = 0$. Take any morphism

$$\begin{array}{ccccccc} T & & 0 & \longrightarrow & P(eA(1-e)A) \oplus (1-e)A & \xrightarrow{p_e} & eA \\ \downarrow & & \downarrow & & \downarrow f & & \downarrow \\ T'[1] & & P(e'A(1-e')A) \oplus (1-e')A & \xrightarrow{p_{e'}} & e'A & \longrightarrow & 0 \end{array}$$

Since $P(eA(1-e)A) \oplus (1-e)A \in \mathsf{add}(1-e)A \subset \mathsf{add}(1-e')A$ and $p_{e'}$ is a right $\mathsf{add}(1-e')A$-approximation, f factors through $p_{e'}$. Thus we have $\mathrm{Hom}_{\mathsf{K}^{\mathrm{b}}(\text{proj-}A)}(T, T'[1]) = 0$, so $T \geq T'$.

Assume $T \geq T'$. Then we have $\mathrm{Hom}_{\mathsf{K}^{\mathrm{b}}(\text{proj-}A)}(T, T'[1]) = 0$. In particular, for any morphism $f : P(eA(1-e)A) \oplus (1-e)A \to e'A$, there exist s and t such that $f = p_{e'}s + tp_e$. Since both p_e and $p_{e'}$ belongs to $J_{\text{mod-}A}$, any morphism in $\mathrm{Hom}_A(P(eA(1-e)A) \oplus (1-e)A, e'A)$ belongs to $J_{\text{mod-}A}$. In particular $(1-e)A$ and $e'A$ do not have a non-zero common summand, and we have $\mathsf{add}eA \supseteq \mathsf{add}e'A$. □

We end this section by observing a connection between BB and APR tilting modules and silting mutation.

Definition 2.52. Let $e \in A$ be a primitive idempotent and S the corresponding simple A-module. Assume $\mathrm{Ext}^1_A(S, S) = 0$, $\mathrm{pd}(\tau^{-1}S) \leq 1$ and that S is not injective. We define a *BB tilting module* [BB] with respect to e by

$$T := \tau^{-1}S \oplus (1-e)A$$

A BB tilting module is called an *APR tilting module* [APR] if S is projective.

Theorem 2.53. *Let T be the BB tilting module with respect to e.*

(a) *T is isomorphic to left mutation $\mu_{eA}^{+}(A)$ of A with respect to eA.*
(b) *T is a tilting A-module of projective dimension at most one.*

Proof. (a) Take a minimal injective resolution of the A-module S:

$$0 \longrightarrow S \longrightarrow D(Ae) \xrightarrow{f} I$$

Since $\mathrm{Ext}^1_A(S,S)=0$, we have $I \in \mathsf{add}D(A(1-e))$. Thus f is a minimal left $\mathsf{add}D(A(1-e))$-approximation. Applying ν^{-1}, we have an exact sequence

$$eA \xrightarrow{\nu^{-1}f} \nu^{-1}I \longrightarrow \tau^{-1}S \longrightarrow 0$$

with a minimal left $(1-e)A$-approximation $\nu^{-1}f$. Since $\mathrm{pd}(\tau^{-1}S) \leq 1$ and S is not injective, we have that $\nu^{-1}f$ is injective. Thus we have $T \simeq \nu^+_{eA}(A)$.

(b) Since $\nu^{-1}f$ is injective, so is $\mathrm{Hom}_{\mathsf{K}^{\mathrm{b}}(\text{proj-}A)}((1-e)A, \nu^{-1}f)$. Thus the assertion follows from Theorem 2.32. □

3. Transitivity for piecewise hereditary algebras

The aim of this section is to prove the following result.

Theorem 3.1. *Let $\mathcal{T} = \mathsf{K}^{\mathrm{b}}(\text{proj-}A)$ where A is either a hereditary algebra or a canonical algebra over a field k. Then the action of iterated irreducible silting mutation on $\mathrm{silt}\,\mathcal{T}$ is transitive.*

The idea of our proof of Theorem 3.1 is to compare silting objects with exceptional sequences.

In the rest of this section let $\mathcal{T}$ be a triangulated category satisfying the following property.

Assumption 3.2. For any $X, Y \in \mathcal{T}$ we have $\mathrm{Hom}_{\mathcal{T}}(X, Y[\ell]) = 0$ for any $|\ell| \gg 0$.

Definition 3.3. Let $\mathcal{T}$ be a triangulated category. We say that an object $X \in \mathcal{T}$ is *exceptional* if $\mathrm{Hom}_{\mathcal{T}}(X, X[\neq 0]) = 0$ and $\mathrm{End}_{\mathcal{T}}(X)$ is a division algebra.

We say that a sequence $(X_1, \cdots, X_n)$ of exceptional objects in $\mathcal{T}$ is an *exceptional sequence* if

$$\mathrm{Hom}_{\mathcal{T}}(X_i, X_j[\mathbb{Z}]) = 0 \quad \text{for any} \quad 1 \leq j < i \leq n.$$

We say that an exceptional sequence is *full* if $\mathsf{thick}(\coprod_{i=1}^n X_i) = \mathcal{T}$. We denote by $\exp\mathcal{T}$ the set of isomorphism classes of full exceptional sequences in $\mathcal{T}$.

Clearly $\mathbb{Z}^n$ acts on $\exp\mathcal{T}$ by

$$(\ell_1, \cdots, \ell_n)(X_1, \cdots, X_n) := (X_1[\ell_1], \cdots, X_n[\ell_n]).$$

Let B_n be the *braid group* generated by $\sigma_1, \cdots, \sigma_{n-1}$ with relations

$$\sigma_i\sigma_j = \sigma_j\sigma_i \quad |i-j| > 1,$$
$$\sigma_i\sigma_{i+1}\sigma_i = \sigma_{i+1}\sigma_i\sigma_{i+1}.$$

Then B_n acts on $\exp\mathcal{T}$ [GR] as follows: For an exceptional sequence $\mathbf{X} := (X_1, \cdots, X_n)$ and $1 \leq i < n$, define objects $L_{X_{i+1}}X_i$ and $R_{X_i}X_{i+1}$ in $\mathcal{T}$ by

$$X_i \longrightarrow \coprod_{\ell\in\mathbb{Z}} D\,\mathrm{Hom}_{\mathcal{T}}(X_i, X_{i+1}[\ell]) \otimes_k X_{i+1}[\ell] \longrightarrow L_{X_{i+1}}X_i[1] \longrightarrow X_i[1]$$

$$X_{i+1}[-1] \longrightarrow R_{X_i}X_{i+1} \longrightarrow \coprod_{\ell\in\mathbb{Z}} \mathrm{Hom}_{\mathcal{T}}(X_i[\ell], X_{i+1}) \otimes_k X_i[\ell] \longrightarrow X_{i+1}$$

and put

$$\begin{aligned} \sigma_i^{-1}\mathbf{X} &:= (X_1, \cdots, X_{i-1}, X_{i+1}, L_{X_{i+1}}X_i, X_{i+2}, \cdots, X_n), \\ \sigma_i\mathbf{X} &:= (X_1, \cdots, X_{i-1}, R_{X_i}X_{i+1}, X_{i+1}, X_{i+2}, \cdots, X_n). \end{aligned}$$

The following transitivity result of exceptional sequences is well-known.

Theorem 3.4. [C, Rin, KM] *Let $\mathcal{T} = \mathsf{K}^{\mathrm{b}}(\text{proj-}A)$ where A is either a hereditary algebra or a canonical algebra over a field k. Then $B_n \times \mathbb{Z}^n$ acts on $\exp\mathcal{T}$ transitively.*

We shall deduce Theorem 3.1 from Theorem 3.4.

We have the following connection between silting objects and exceptional sequences, asserting that any full exceptional sequence gives rise to a silting object.

Proposition 3.5. *Let $\mathbf{X} = (X_1, \cdots, X_n)$ be a full exceptional sequence in $\mathcal{T}$. Then there exists $a \in \mathbb{Z}$ such that $X_1[\ell_1] \oplus \cdots \oplus X_n[\ell_n]$ is a silting object for any integers $\ell_1 \cdots, \ell_n \in \mathbb{Z}$ satisfying $\ell_i + a \leq \ell_{i+1}$ for any $1 \leq i < n$.*

Proof. By Assumption 3.2, there exists $a \geq 0$ such that $\mathrm{Hom}_{\mathcal{T}}(X_i, X_j[> a]) = 0$ and any $1 \leq i, j \leq n$. We shall show that a satisfies the desired condition.

Let $\ell_1 \cdots, \ell_n$ be integers satisfying $\ell_i + a \leq \ell_{i+1}$ for any i. Fix $1 \leq i < j \leq n$. Since $\mathbf{X} \in \exp \mathcal{T}$, we have $\mathrm{Hom}_{\mathcal{T}}(X_j[\ell_j], X_i[\ell_i + \ell]) = 0$ for any $\ell \in \mathbb{Z}$ and $\mathrm{Hom}_{\mathcal{T}}(X_i[\ell_i], X_i[\ell_i + \ell]) = 0$ for any $\ell \neq 0$. On the other hand we have $\mathrm{Hom}_{\mathcal{T}}(X_i[\ell_i], X_j[\ell_j + \ell]) = 0$ for any $\ell > 0$ by $\ell_j + \ell - \ell_i > a$ and our choice of a. Thus we have the assertion. □

We have the following easy observation.

Lemma 3.6. *Let $\mathbf{X} = (X_1, \cdots, X_n)$ be a full exceptional sequence in $\mathcal{T}$ such that $M = X_1 \oplus \cdots \oplus X_n$ is a silting object. For any integers $\ell_1 \leq \cdots \leq \ell_n$ we have the following.*

1. *$X_1[\ell_1] \oplus \cdots \oplus X_n[\ell_n]$ is a silting object.*
2. *If $\ell_1 \geq 0$, then $X_1 \oplus \cdots \oplus X_n$ and $X_1[\ell_1] \oplus \cdots \oplus X_n[\ell_n]$ are transitive under iterated irreducible silting mutation.*

Proof. (1) Fix $i \leq j$. Since $\mathbf{X} \in \exp \mathcal{T}$, we have $\mathrm{Hom}_{\mathcal{T}}(X_j[\ell_j], X_i[\ell_i + \ell]) = 0$ for any $\ell \neq 0$. Since $X_1 \oplus \cdots \oplus X_n$ is a silting object and $\ell_i \leq \ell_j$, we have $\mathrm{Hom}_{\mathcal{T}}(X_i[\ell_i], X_j[\ell_j + \ell]) = 0$ for any $\ell > 0$. Thus we have the assertion.

(2) We use the induction on $\ell_1 + \cdots + \ell_n$. Let $\ell_0 := 0$. Assume $\ell_{i-1} < \ell_i$ for some $1 \leq i \leq n$. Then $\mathrm{Hom}_{\mathcal{T}}(X_j[\ell_j], X_i[\ell_i]) = 0$ for any $j > i$ since $\mathbf{X} \in \exp \mathcal{T}$, and also for any $j < i$ since $M \in \mathrm{silt}\, \mathcal{T}$ and $\ell_i - \ell_j > 0$. Consequently we have $\mathrm{Hom}_{\mathcal{T}}(X_j[\ell_j], X_i[\ell_i]) = 0$ for any $j \neq i$. Thus we have

$$\mu^-_{X_i[\ell_i]}(X_1[\ell_1] \oplus \cdots \oplus X_n[\ell_n]) = X_1[\ell_1] \oplus \cdots X_i[\ell_i - 1] \cdots \oplus X_n[\ell_n].$$

Thus the assertion follows inductively. □

We have the following transitivity result.

Proposition 3.7. *Let $(X_1, \cdots, X_n)$ be a full exceptional sequence in $\mathcal{T}$ and $\ell_1, \cdots, \ell_n \in \mathbb{Z}$, If $X_1 \oplus \cdots \oplus X_n$ and $X_1[\ell_1] \oplus \cdots \oplus X_n[\ell_n]$ are silting objects, then they are transitive under iterated irreducible silting mutation.*

Proof. Choose integers $0 \leq m_1 \leq \cdots \leq m_n$ satisfying $0 \leq m_1 - \ell_1 \leq \cdots \leq m_n - \ell_n$. By Lemma 3.6 we have that $X_1[m_1] \oplus \cdots \oplus X_n[m_n]$ is an iterated irreducible silting mutation of $X_1 \oplus \cdots \oplus X_n$ (respectively, $X_1[\ell_1] \oplus \cdots \oplus X_n[\ell_n]$). Thus we have the assertion. □

In the rest of this section let $\mathcal{T} = \mathsf{K}^{\mathrm{b}}(\text{proj-}A)$ where A is either a hereditary algebra or a canonical algebra over a field. We can identify $\mathcal{T}$ with $\mathsf{D}^{\mathrm{b}}(\mathcal{A})$ for a hereditary abelian category $\mathcal{A}$. Any indecomposable object in $\mathcal{T}$ is isomorphic to $X[\ell]$ for some $X \in \mathcal{A}$ and $\ell \in \mathbb{Z}$. Moreover $\mathcal{T}$ satisfies Assumption 3.2.

Let us start with the following preliminary results.

Lemma 3.8. [H, IV.1.2, IV.1.5]

1. *Let $X, Y \in \mathcal{A}$ be indecomposable objects satisfying $\mathrm{Ext}^1_{\mathcal{A}}(Y, X) = 0$. Then any non-zero morphism $f \in \mathrm{Hom}_{\mathcal{A}}(X, Y)$ is either injective or surjective.*
2. *Let $X, Y \in \mathcal{A}$ be exceptional objects. If $\mathrm{Hom}_{\mathcal{A}}(X, Y) \neq 0$ and $\mathrm{Ext}^1_{\mathcal{A}}(X, Y) \neq 0$, then we have $\mathrm{Ext}^1_{\mathcal{A}}(Y, X) \neq 0$.*

We have the following result asserting that $\mathbf{R}\,\mathrm{Hom}_{\mathcal{A}}(X, Y)$ is a stalk complex.

Lemma 3.9. *Let $X, Y \in \mathcal{T}$ be indecomposable objects. If $M := X \oplus Y$ satisfies $\mathrm{Hom}_{\mathcal{T}}(M, M[> 0]) = 0$, then there exists at most one $\ell \in \mathbb{Z}$ such that $\mathrm{Hom}_{\mathcal{T}}(X, Y[\ell]) \neq 0$.*

Proof. Without loss of generality, we can assume that X and $Z := Y[a]$ belong to $\mathcal{A}$. Since $\mathcal{A}$ is hereditary, we have $\mathrm{Hom}_{\mathcal{T}}(X, Z[\neq 0, 1]) = 0$. Assume that $\mathrm{Hom}_{\mathcal{T}}(X, Z) \neq 0$ and $\mathrm{Hom}_{\mathcal{T}}(X, Z[1]) \neq 0$. By Lemma 3.8(2) we have $\mathrm{Hom}_{\mathcal{T}}(Z, X[1]) \neq 0$.

Since $\mathrm{Hom}_{\mathcal{T}}(X, Z[1]) \neq 0$, we have $\mathrm{Hom}_{\mathcal{T}}(M, M[1+a]) \neq 0$. Thus we have $a < 0$. On the other hand, since $\mathrm{Hom}_{\mathcal{T}}(Z, X[1]) \neq 0$, we have $\mathrm{Hom}_{\mathcal{T}}(M[a], M[1]) \neq 0$. Thus we have $a > 0$, a contradiction. □

We have the following information about existence of ℓ in Lemma 3.9.

Lemma 3.10. *Let $X_{n+1} = X_1, \cdots, X_n \in \mathcal{T}$ be pairwise non-isomorphic indecomposable objects. Assume that the following conditions are satisfied.*

(i) *$M := X_1 \oplus \cdots \oplus X_n$ satisfies $\mathrm{Hom}_{\mathcal{T}}(M, M[>0]) = 0$.*
(ii) *There exist integers $\ell_1, \cdots, \ell_n \in \mathbb{Z}$ such that $\mathrm{Hom}_{\mathcal{T}}(X_i, X_{i+1}[\ell_i]) \neq 0$ for any $1 \leq i \leq n$.*

Then we have $n = 1$.

Proof. By (i) we have $\ell_i \leq 0$ for any i. We take $a_i \in \mathbb{Z}$ such that $X_i \in \mathcal{A}[a_i]$. Since $\mathrm{Hom}_{\mathcal{T}}(X_i, X_{i+1}[\ell_i]) \neq 0$, we have $\mathrm{Hom}_{\mathcal{T}}(\mathcal{A}[a_i], \mathcal{A}[a_{i+1} + \ell_i]) \neq 0$. Thus $a_{i+1} - a_i + \ell_i$ is either 0 or 1. We have

$$0 \geq \sum_{i=1}^{n} \ell_i = \sum_{i=1}^{n} (a_{i+1} - a_i + \ell_i) = \sum_{i=1}^{n} (0 \text{ or } 1),$$

which implies $\ell_1 = \cdots = \ell_n = 0$ and $a_1 = \cdots = a_n$. Thus we can assume that each X_i belongs to $\mathcal{A}$.

Assume $n > 1$ and take a non-zero morphism $f_i : X_i \to X_{i+1}$ for any $1 \leq i \leq n$. Let $f_{n+1} := f_1$. Then f_i is not an isomorphism and either injective or surjective by Proposition 3.8(1). Also each $f_{i+1} f_i$ is either injective or surjective again by Proposition 3.8(1). So it is impossible that f_i is surjective and f_{i+1} is injective at the same time. It is easy to conclude that either all f_i are injective or all f_i are surjective. Thus $f_n \cdots f_1 : X_1 \to X_1$ is not an isomorphism and either injective or surjective. This is a contradiction. □

Now we show a certain converse of Proposition 3.5 asserting that any silting object gives rise to a full exceptional sequence. It is also possible to show this observation by applying [AST, Theorem A].

Proposition 3.11. *Let $M = X_1 \oplus \cdots \oplus X_n$ be a basic silting object in $\mathcal{T}$. Then we can change indices of $X_1, \cdots, X_n$ such that $(X_1, \cdots, X_n)$ is a full exceptional sequence in $\mathcal{T}$.*

Proof. For each $1 \leq i, j \leq n$, we write $X_i \leq X_j$ if there exists a sequence $X_i = X_{i_1}, X_{i_2}, \cdots, X_{i_m} = X_j$ such that $\mathrm{Hom}_{\mathcal{T}}(X_{i_a}, X_{i_{a+1}}[\ell_a]) \neq 0$ for some $\ell_a \in \mathbb{Z}$. Clearly we have $X_i \leq X_i$ and that $X_i \leq X_j \leq X_k$ implies $X_i \leq X_k$. By Lemma 3.10 we have that $X_i \leq X_j \leq X_i$ implies $i = j$. This means that $X_1, \cdots, X_n$ forms a partially ordered set. Thus we can change indices of $X_1, \cdots, X_n$ such that $X_i \leq X_j$ implies $i \leq j$. Then $(X_1, \cdots, X_n)$ forms an exceptional sequence. □

For an exceptional sequence $\mathbf{X} = (X_1, \cdots, X_n)$, we let

$$[[\mathbf{X}]] := X_1 \oplus \cdots \oplus X_n \in \mathcal{T}.$$

The following is a main inductive step in the proof of Theorem 3.1.

Lemma 3.12. *Let $\mathbf{X} = (X_1, \cdots, X_n)$ be a full exceptional sequence in $\mathcal{T}$ such that $[[\mathbf{X}]]$ is a silting object. For any $1 \leq i < n$, there exists $\boldsymbol{\ell} \in \mathbb{Z}^n$ (respectively, $\boldsymbol{m} \in \mathbb{Z}^n$) such that $[[(\sigma_i, \boldsymbol{\ell})\mathbf{X}]]$ (respectively, $[[(\sigma_i^{-1}, \boldsymbol{m})\mathbf{X}]]$) is a iterated irreducible silting mutation of $[[\mathbf{X}]]$.*

Proof. By Lemma 3.9 there exists at most one $a \in \mathbb{Z}$ such that $\mathrm{Hom}_{\mathcal{T}}(X_i, X_{i+1}[a]) \neq 0$. Then a must be non-positive since $[[\mathbf{X}]]$ is silting. If such a does not exist, we let $a := 0$.

By Assumption 3.2 there exists $b \leq 0$ such that

$$\mathrm{Hom}_{\mathcal{T}}(X_j[b], X_{i+1}[\ell]) = 0 \quad \text{for any } \ 1 \leq j < i \ \text{ and } \ a \leq \ell \leq 0. \tag{9}$$

By Lemma 3.6 we have an exceptional sequence

$$\mathbf{Y} := (X_1[b], \cdots, X_{i-1}[b], X_i, X_{i+1}, X_{i+2}, \cdots, X_n)$$

such that $[[\mathbf{Y}]]$ is an iterated irreducible silting mutation of $[[\mathbf{X}]]$. Let

$$\begin{aligned}\mathbf{Z} &:= (X_1[b], \cdots, X_{i-1}[b], X_i, X_{i+1}[a], X_{i+2}, \cdots, X_n),\\ \sigma_i\mathbf{Z} &= (X_1[b], \cdots, X_{i-1}[b], R_{X_i}(X_{i+1}[a]), X_i, X_{i+2}, \cdots, X_n).\end{aligned}$$

By our choice of a and (9) we have

$$\mu_{i+1}^{-(-a)}([[\mathbf{Y}]]) = [[\mathbf{Z}]] \text{ and } \mu_{i+1}^{-(-a+1)}([[\mathbf{Y}]]) = [[\sigma_i\mathbf{Z}]].$$

Consequently $[[\sigma_i\mathbf{Z}]] = [[(\sigma_i, \boldsymbol{\ell})\mathbf{X}]]$ is an iterated irreducible silting mutation of $[[\mathbf{X}]]$, where we put $\boldsymbol{\ell} := (b, \cdots, b, 0, a, 0, \cdots, 0)$. □

Now we are ready to prove Theorem 3.1.

Let $T = X_1 \oplus \cdots \oplus X_n$ and $U = Y_1 \oplus \cdots \oplus Y_n$ be basic silting objects in $\mathcal{T}$. By Proposition 3.11 we can assume that $\mathbf{X} = (X_1, \cdots, X_n)$ and $\mathbf{Y} = (Y_1, \cdots, Y_n)$ are full exceptional sequences. By Theorem 3.4 there exists $(\sigma, \boldsymbol{\ell}) \in B_n \times \mathbb{Z}^n$ such that $\mathbf{Y} = (\sigma, \boldsymbol{\ell})\mathbf{X}$. Writing down $\sigma \in B_n$ as a product of $\sigma_1^{\pm 1}, \cdots, \sigma_{n-1}^{\pm 1}$ and applying Lemma 3.12 repeatedly, we have $\boldsymbol{m} \in \mathbb{Z}^n$ such that $[[(\sigma, \boldsymbol{m})\mathbf{X}]]$ is an iterated irreducible silting mutation of $[[\mathbf{X}]]$. By Proposition 3.7 $[[(\sigma, \boldsymbol{m})\mathbf{X}]] = [[(0, \boldsymbol{m} - \boldsymbol{\ell})\mathbf{Y}]]$ is an iterated irreducible silting mutation of $[[\mathbf{Y}]]$. Thus we have the assertion. □

4. Silting subcategories in triangulated categories with coproducts

The aim of this section is to study silting subcategories for triangulated categories which have arbitrary coproducts. We need to modify the definition of silting subcategories (Definition 4.1) from that in section 2. The advantage of this setting is that each set of compact objects gives rise to a torsion pair (Theorem 4.3), which is not the case for the setting of section 2. As an application, we deduce a result of Hoshino-Kato-Miyachi [HKM], which associates a t-structure for each silting subcategory (Corollary 4.7). Moreover we show that this gives a one-to-one correspondence between silting subcategories and certain t-structures (Theorem 4.10). We also deduce a result of Paukszello [P1, P2] on co-t-structures.

Throughout this section, let $\mathcal{T}$ be a triangulated category with arbitrary coproducts.

We say that an object $X \in \mathcal{T}$ is *compact* if $\mathrm{Hom}_{\mathcal{T}}(X, -)$ commutes with arbitrary coproducts. We denote by $\mathcal{T}^{\mathrm{c}}$ the full subcategory consisting of compact objects in $\mathcal{T}$. We say that a subcategory $\mathcal{M}$ of $\mathcal{T}$ is *compact* if $\mathcal{M} \subset \mathcal{T}^{\mathrm{c}}$, *generating* if $\mathcal{M}[\mathbb{Z}]^{\perp_{\mathcal{T}}} = 0$, and *skeletally small* if isomorphism classes of objects in $\mathcal{M}$ form a set.

Definition 4.1. We say that a subcategory $\mathcal{M}$ of $\mathcal{T}$ is *silting* if $\mathrm{Hom}_{\mathcal{T}}(\mathcal{M}, \mathcal{M}[>0]) = 0$ and $\mathcal{M}$ is a skeletally small compact and generating subcategory of $\mathcal{T}$.

In this case $\mathcal{M}$ is a silting subcategory of $\mathcal{T}^{\mathrm{c}}$ in the sense of Definition 2.1 by the following Neeman's result:

Proposition 4.2. [N] *For any compact generating subcategory $\mathcal{M}$ of $\mathcal{T}$, we have* $\mathsf{thick}\mathcal{M} = \mathcal{T}^{\mathrm{c}}$.

4.1. **Torsion pairs induced by sets of compact objects.** The following main result in this section asserts that each set of compact objects gives a t-structure.

Theorem 4.3. *Let $\mathcal{T}$ be a triangulated category with arbitrary coproducts, and let $\mathcal{C}$ be a set of objects in $\mathcal{T}^{\mathrm{c}}$. Then $({}^{\perp}(\mathcal{C}^{\perp}), \mathcal{C}^{\perp})$ is a torsion pair.*

We need the following general observation of homotopy colimit [N].

Proposition 4.4. *Let $\mathcal{T}$ be a triangulated category with arbitrary coproducts. For a sequence*

$$X_0 \xrightarrow{a_0} X_1 \xrightarrow{a_1} X_2 \xrightarrow{a_2} \cdots$$

we put

$$1 - \mathrm{shift} := \begin{pmatrix} 1 & 0 & 0 & 0 & \cdots \\ -a_0 & 1 & 0 & 0 & \cdots \\ 0 & -a_1 & 1 & 0 & \cdots \\ 0 & 0 & -a_2 & 1 & \cdots \\ \vdots & \vdots & \vdots & \vdots & \ddots \end{pmatrix} : \coprod_{i \geq 0} X_i \to \coprod_{i \geq 0} X_i$$

and take a triangle

$$\coprod_{i\geq 0} X_i \xrightarrow{1-\text{shift}} \coprod_{i\geq 0} X_i \to Y \to \coprod_{i\geq 0} X_i[1],$$

where Y is called the homotopy colimit. *Then we have the following isomorphism on $\mathcal{T}^{\mathrm{c}}$:*

$$\mathrm{Hom}_{\mathcal{T}}(-,Y) \simeq \mathrm{colim}(\mathrm{Hom}_{\mathcal{T}}(-,X_0) \xrightarrow{a_0\cdot} \mathrm{Hom}_{\mathcal{T}}(-,X_1) \xrightarrow{a_1\cdot} \mathrm{Hom}_{\mathcal{T}}(-,X_2) \xrightarrow{a_2\cdot} \cdots).$$

We are ready to prove our Theorem 4.3.

For any $X \in \mathcal{T}$, we only have to construct a triangle $S_X \to X \to U_X \to S_X[1]$ in $\mathcal{T}$ with $U_X \in \mathcal{C}^{\perp}$ and $S_X \in {}^{\perp}(\mathcal{C}^{\perp})$. This is given as follows:

Proposition 4.5. *Fix $X = X_0 \in \mathcal{T}$. For each $i \geq 0$ we take a triangle*

$$C_i \xrightarrow{b_i} X_i \xrightarrow{a_i} X_{i+1} \longrightarrow C_i[1] \tag{10}$$

such that b_i is a right $\mathsf{Add}\mathcal{C}$-approximation. This is possible since $\mathcal{C}$ forms a set. Thus we have a sequence

$$\begin{array}{ccccccccc} C_0 & & C_1 & & C_2 & & & & \\ {\scriptstyle b_0}\downarrow & & {\scriptstyle b_1}\downarrow & & {\scriptstyle b_2}\downarrow & & & & \\ X_0 & \xrightarrow{a_0} & X_1 & \xrightarrow{a_1} & X_2 & \xrightarrow{a_2} & \cdots & & \end{array}$$

of morphisms. We denote by ι_i the inclusion $X_i \to \coprod_{i\geq 0} X_i$. We take a triangle

$$\coprod_{i\geq 0} X_i \xrightarrow{1-\text{shift}} \coprod_{i\geq 0} X_i \xrightarrow{c} U_X \to \coprod_{i\geq 0} X_i[1] \tag{11}$$

and for the composition $d := c\iota_0 : X \to U_X$ we take a triangle

$$S_X \longrightarrow X \xrightarrow{d} U_X \longrightarrow S_X[1]. \tag{12}$$

Then we have $S_X \in {}^{\perp}(\mathcal{C}^{\perp})$ and $U_X \in \mathcal{C}^{\perp}$.

Proof. (i) We shall show $U_X \in \mathcal{C}^{\perp}$.

For any $i \geq 0$ we have an exact sequence

$$\mathrm{Hom}_{\mathcal{T}}(\mathcal{C}, C_i) \xrightarrow{b_i\cdot} \mathrm{Hom}_{\mathcal{T}}(\mathcal{C}, X_i) \xrightarrow{a_i\cdot} \mathrm{Hom}_{\mathcal{T}}(\mathcal{C}, X_{i+1}).$$

Since b_i is a right $\mathsf{Add}\mathcal{C}$-approximation, the left map is surjective, and hence the right map is zero. Since $\mathcal{C}$ consists of compact objects, we have by Proposition 4.4

$$\mathrm{Hom}_{\mathcal{T}}(\mathcal{C}, U_X) \simeq \mathrm{colim}(\mathrm{Hom}_{\mathcal{T}}(\mathcal{C}, X_0) \xrightarrow{0} \mathrm{Hom}_{\mathcal{T}}(\mathcal{C}, X_1) \xrightarrow{0} \cdots) = 0.$$

Thus we have $U_X \in \mathcal{C}^{\perp}$.

(ii) We shall show that $d : X \to U_X$ is a left $\mathcal{C}^{\perp}$-approximation.

Fix any $f_0 : X = X_0 \to Y$ with $Y \in \mathcal{C}^{\perp}$. Since $f_0 b_0 \in \mathrm{Hom}_{\mathcal{T}}(C_0, Y) = 0$, there exists $f_1 : X_1 \to Y$ such that $f_0 = f_1 a_0$ by the triangle (10). Similarly we have $f_i : X_i \to Y$ for any $i > 0$ such that

$$f_{i-1} = f_i a_{i-1}. \tag{13}$$

$$\begin{array}{ccccccccc} C_0 & & C_1 & & C_2 & & & & \\ {\scriptstyle b_0}\downarrow & & {\scriptstyle b_1}\downarrow & & {\scriptstyle b_2}\downarrow & & & & \\ X_0 & \xrightarrow{a_0} & X_1 & \xrightarrow{a_1} & X_2 & \xrightarrow{a_2} & \cdots & & \\ {\scriptstyle f_0}\downarrow & \swarrow_{f_1} & & \swarrow_{f_2} & & & & & \\ Y & & & & & & & & \end{array}$$

Now we consider a morphism

$$f := (f_0\ f_1\ \cdots) \in \mathrm{Hom}_{\mathcal{T}}(\coprod_{i\geq 0} X_i, Y).$$

By (13) we have $f(1-\text{shift}) = 0$. By the triangle (11) there exists $g \in \mathrm{Hom}_{\mathcal{T}}(U_X, Y)$ such that $f = gc$. In particular we have $f_0 = f\iota_0 = gd$.

(iii) We shall show that $\mathrm{Hom}_{\mathcal{T}}(\coprod_{i\geq 0} X_i, \mathcal{C}^{\perp}) \xrightarrow{\cdot(1-\mathrm{shift})} \mathrm{Hom}_{\mathcal{T}}(\coprod_{i\geq 0} X_i, \mathcal{C}^{\perp})$ is surjective.

Let $Y \in \mathcal{C}^{\perp}$. Fix any $f = (f_0\ f_1\ \cdots) \in \mathrm{Hom}_{\mathcal{T}}(\coprod_{i\geq 0} X_i, Y)$. Since $f_0 b_0 = 0$, there exists $g_1 \in \mathrm{Hom}_{\mathcal{T}}(X_1, Y)$ such that $f_0 = -g_1 a_0$ by the triangle (10). Since $(f_1 - g_1)b_1 = 0$, there exists $g_2 \in \mathrm{Hom}_{\mathcal{T}}(X_2, Y)$ such that $f_1 - g_1 = -g_2 a_1$. Similarly we have $g_i : X_i \to Y$ for any $i \geq 0$ which makes the following diagram commutative.

$$\begin{array}{ccccccc} C_i & \xrightarrow{b_i} & X_i & \xrightarrow{a_i} & X_{i+1} & \longrightarrow & C_i[1] \\ & & {\scriptstyle f_i - g_i}\downarrow & \swarrow{\scriptstyle -g_{i+1}} & & & \\ & & Y & & & & \end{array}$$

Now the morphism $g := (0\ g_1\ g_2\ \cdots)$ satisfies $f = g(1 - \mathrm{shift})$.

(iv) We shall show that $\mathrm{Hom}_{\mathcal{T}}(U_X, \mathcal{C}^{\perp}[1]) \xrightarrow{\cdot d} \mathrm{Hom}_{\mathcal{T}}(X, \mathcal{C}^{\perp}[1])$ is injective.

Let $Y \in \mathcal{C}^{\perp}[1]$. Assume that $f \in \mathrm{Hom}_{\mathcal{T}}(U_X, Y)$ satisfies $fd = 0$. Put $g = (g_0\ g_1\ \cdots) := fc \in \mathrm{Hom}_{\mathcal{T}}(\coprod_{i\geq 0} X_i, Y)$. Then we have $g_0 = fc\iota_0 = fd = 0$.

$$\begin{array}{ccccccccc} & & & & & & Y & & \\ & & & & & {\scriptstyle g}\nearrow & {\scriptstyle f}\uparrow & & \\ \coprod_{i\geq 0} X_i & \xrightarrow{1-\mathrm{shift}} & \coprod_{i\geq 0} X_i & & \xrightarrow{c} & & U_X & \longrightarrow & \coprod_{i\geq 0} X_i[1] \\ \uparrow & & \uparrow{\scriptstyle \iota_0} & & & & \| & & \uparrow \\ S_X & \longrightarrow & X & & \xrightarrow{d} & & U_X & \longrightarrow & S_X[1] \end{array}$$

Assume $g_i = 0$ for some $i \geq 0$. Since

$$-g_{i+1}a_i = g_i - g_{i+1}a_i = g(1 - \mathrm{shift})\iota_i = fc(1 - \mathrm{shift})\iota_i = 0,$$

we have that g_{i+1} factors through $C_i[1]$ by the triangle (10). Since $Y \in \mathcal{C}^{\perp}[1]$, we have $g_{i+1} = 0$.

Inductively we have $g = 0$, so f factors through $U_X \to \coprod_{i\geq 0} X_i[1]$. Now using (iii) we have $f = 0$.

(v) We shall show $S_X \in {}^{\perp}(\mathcal{C}^{\perp})$.

We have an exact sequence

$$\mathrm{Hom}_{\mathcal{T}}(U_X, \mathcal{C}^{\perp}) \xrightarrow{\cdot d} \mathrm{Hom}_{\mathcal{T}}(X, \mathcal{C}^{\perp}) \to \mathrm{Hom}_{\mathcal{T}}(S_X, \mathcal{C}^{\perp}) \to \mathrm{Hom}_{\mathcal{T}}(U_X[-1], \mathcal{C}^{\perp}) \xrightarrow{\cdot d[-1]} \mathrm{Hom}_{\mathcal{T}}(X[-1], \mathcal{C}^{\perp}).$$

By (ii) the left map $(\cdot d)$ is surjective, and by (iv) the right map $(\cdot d[-1])$ is injective. Thus we have $\mathrm{Hom}_{\mathcal{C}}(S_X, \mathcal{C}^{\perp}) = 0$, and we have the assertion. □

As special cases of Theorem 4.3, we have the following results due to Beligiannis-Reiten and Pauksztello.

Corollary 4.6. *Let $\mathcal{T}$ be a triangulated category with arbitrary coproducts, and $\mathcal{C}$ a set of objects in $\mathcal{T}^{\mathrm{c}}$.*

(a) [BR] *If $\mathcal{C}[1] \subset \mathcal{C}$, then $({}^{\perp}(\mathcal{C}^{\perp}), \mathcal{C}^{\perp}[1])$ is a t-structure.*
(b) [P2] *If $\mathcal{C} \subset \mathcal{C}[1]$, then $({}^{\perp}(\mathcal{C}^{\perp})[1], \mathcal{C}^{\perp})$ is a co-t-structure.*

Let us apply our results to more special cases, where we can describe the category ${}^{\perp}(\mathcal{C}^{\perp})$ in a more direct way.

The first application is the following result (b) of Hoshino-Kato-Miyachi [HKM].

Corollary 4.7. *Let $\mathcal{T}$ be a triangulated category with arbitrary coproducts and $\mathcal{M}$ a skeletally small compact subcategory satisfying* $\mathrm{Hom}_{\mathcal{T}}(\mathcal{M}, \mathcal{M}[>0]) = 0$.

(a) *We have ${}^{\perp}(\mathcal{M}[>0]^{\perp}) \subseteq \mathcal{M}[\leq 0]^{\perp}$.*
(b) *If $\mathcal{M}$ is a silting subcategory, then equality in (a) holds and $(\mathcal{M}[<0]^{\perp}, \mathcal{M}[>0]^{\perp})$ is a t-structure of $\mathcal{T}$.*

Proof. Let $\mathcal{C} := \mathcal{M}[>0]$. For any $X \in \mathcal{T}$, we apply Proposition 4.5 to get a triangle $S_X \to X \xrightarrow{d} U_X \to S_X[1]$ with $S_X \in {}^{\perp}(\mathcal{C}^{\perp})$ and $U_X \in \mathcal{C}^{\perp}$.

Applying $\mathrm{Hom}_{\mathcal{T}}(\mathcal{M}[\leq 0], -)$ to (10) for any $i \geq 0$, we have an isomorphism

$$(a_i \cdot) : \mathrm{Hom}_{\mathcal{T}}(\mathcal{M}[\leq 0], X_i) \xrightarrow{\sim} \mathrm{Hom}_{\mathcal{T}}(\mathcal{M}[\leq 0], X_{i+1}).$$

Since $\mathcal{M}[\leq 0]$ is compact, Proposition 4.4 gives an isomorphism

$$(d\cdot) : \mathrm{Hom}_{\mathcal{T}}(\mathcal{M}[\leq 0], X) \xrightarrow{\sim} \mathrm{Hom}_{\mathcal{T}}(\mathcal{M}[\leq 0], U_X). \tag{14}$$

(a) Let $X \in {}^{\perp}(\mathcal{C}^{\perp})$. Then the morphism $d : X \to U_X$ is zero since $U_X \in \mathcal{C}^{\perp}$. Since (14) is an isomorphism which is zero, we have $\mathrm{Hom}_{\mathcal{T}}(\mathcal{M}[\leq 0], X) = 0$.

(b) By Theorem 4.3 we only have to show ${}^{\perp}(\mathcal{C}^{\perp}) = \mathcal{M}[\leq 0]^{\perp}$. By (a), we only have to show ${}^{\perp}(\mathcal{C}^{\perp}) \supset \mathcal{M}[\leq 0]^{\perp}$. Let $X \in \mathcal{M}[\leq 0]^{\perp}$. By (14), we have $\mathrm{Hom}_{\mathcal{T}}(\mathcal{M}[\leq 0], U_X) \simeq \mathrm{Hom}_{\mathcal{T}}(\mathcal{M}[\leq 0], X) = 0$. Thus $U_X \in \mathcal{C}^{\perp} \cap \mathcal{M}[\leq 0]^{\perp} = \mathcal{M}[\mathbb{Z}]^{\perp} = 0$, and we have $X \simeq S_X \in {}^{\perp}(\mathcal{C}^{\perp})$. □

The second application is Corollary 4.8 below, which is a version of a result of Pauksztello [P1] on co-t-structures, where the assumption $\mathrm{Hom}_{\mathcal{T}}(\mathcal{M}, \mathcal{M}[1]) = 0$ in [P1] is dropped.

We say that a subcategory $\mathcal{M}$ of $\mathcal{T}$ is *cosilting* if $\mathrm{Hom}_{\mathcal{T}}(\mathcal{M}, \mathcal{M}[< 0]) = 0$ and $\mathcal{M}$ is a skeletally small compact and generating subcategory of $\mathcal{T}$.

We say that an additive category $\mathcal{M}$ is *semisimple* if $\mathrm{End}_{\mathcal{M}}(X)$ is a semisimple ring for any $X \in \mathcal{M}$.

Corollary 4.8. *Let $\mathcal{T}$ be a triangulated category with arbitrary coproducts and $\mathcal{M}$ a skeletally small compact and semisimple subcategory satisfying* $\mathrm{Hom}_{\mathcal{T}}(\mathcal{M}, \mathcal{M}[< 0]) = 0$.

(a) *We have ${}^{\perp}(\mathcal{M}[< 0]^{\perp}) \subseteq \mathcal{M}[\geq 0]^{\perp}$.*
(b) *If $\mathcal{M}$ is a cosilting subcategory, then equality in (a) holds and $(\mathcal{M}[> 0]^{\perp}, \mathcal{M}[< 0]^{\perp})$ is a co-t-structure of $\mathcal{T}$.*

Proof. Let $\mathcal{C} := \mathcal{M}[< 0]$. For any $X \in \mathcal{T}$, we apply Proposition 4.5 with an additional assumption that $\mathrm{Hom}_{\mathcal{T}}(\mathcal{M}[-1], C_i) \xrightarrow{b_i\cdot} \mathrm{Hom}_{\mathcal{T}}(\mathcal{M}[-1], X_i)$ is an isomorphism for any $i \geq 0$. This is possible since $\mathcal{M}$ is semisimple. Applying $\mathrm{Hom}_{\mathcal{T}}(\mathcal{M}[\geq 0], -)$ to (10) for any $i \geq 0$, we have an isomorphism

$$(a_i\cdot) : \mathrm{Hom}_{\mathcal{T}}(\mathcal{M}[\geq 0], X_i) \xrightarrow{\sim} \mathrm{Hom}_{\mathcal{T}}(\mathcal{M}[\geq 0], X_{i+1}).$$

Since $\mathcal{M}[\geq 0]$ is compact, Proposition 4.4 implies that our triangle $S_X \to X \xrightarrow{d} U_X \to S_X[1]$ induces an isomorphism

$$(d\cdot) : \mathrm{Hom}_{\mathcal{T}}(\mathcal{M}[\geq 0], X) \xrightarrow{\sim} \mathrm{Hom}_{\mathcal{T}}(\mathcal{M}[\geq 0], U_X). \tag{15}$$

(a) Let $X \in {}^{\perp}(\mathcal{C}^{\perp})$. Then the morphism $d : X \to U_X$ is zero since $U_X \in \mathcal{C}^{\perp}$. Since (15) is an isomorphism which is zero, we have $\mathrm{Hom}_{\mathcal{T}}(\mathcal{M}[\geq 0], X) = 0$.

(b) By Theorem 4.3 we only have to show ${}^{\perp}(\mathcal{C}^{\perp}) \supset \mathcal{M}[\geq 0]^{\perp}$. Let $X \in \mathcal{M}[\geq 0]^{\perp}$. By (15), we have $\mathrm{Hom}_{\mathcal{T}}(\mathcal{M}[\geq 0], U_X) \simeq \mathrm{Hom}_{\mathcal{T}}(\mathcal{M}[\geq 0], X) = 0$. Thus $U_X \in \mathcal{C}^{\perp} \cap \mathcal{M}[\geq 0]^{\perp} = \mathcal{M}[\mathbb{Z}]^{\perp} = 0$, and we have $X \simeq S_X \in {}^{\perp}(\mathcal{C}^{\perp})$. □

4.2. **The correspondence between silting subcategories and t-structures.** Let $\mathcal{T}$ be a triangulated category with arbitrary coproducts. For a silting subcategory $\mathcal{M}$ of $\mathcal{T}$, we observed in Corollary 4.7 that we have a t-structure $(\mathcal{T}^{\leq 0}_{\mathcal{M}}, \mathcal{T}^{\geq 0}_{\mathcal{M}})$ given by

$$\mathcal{T}^{\leq 0}_{\mathcal{M}} := {}^{\perp}(\mathcal{M}[\geq 0]^{\perp}) = \mathcal{M}[< 0]^{\perp} \quad \text{and} \quad \mathcal{T}^{\geq 0}_{\mathcal{M}} := \mathcal{M}[> 0]^{\perp}.$$

The aim of this subsection is to characterize these t-structures obtained from silting subcategories. The following definition is suggested by Proposition 2.13.

Definition 4.9. Let $(\mathcal{T}^{\leq 0}, \mathcal{T}^{\geq 0})$ be a t-structure in $\mathcal{T}$ with the heart $\mathcal{T}^0$. Define a subcategory of $\mathcal{T}$ by

$$\mathcal{M} = \mathcal{M}(\mathcal{T}^{\leq 0}, \mathcal{T}^{\geq 0}) := \mathcal{T}^{\leq 0} \cap {}^{\perp}(\mathcal{T}^{< 0}) \cap \mathcal{T}^{\mathrm{c}}.$$

(a) We say that the t-structure $(\mathcal{T}^{\leq 0}, \mathcal{T}^{\geq 0})$ is *silting* if $\mathcal{M}$ is skeletally small and generating.
(b) We say that a silting t-structure is *tilting* if $\mathcal{M} \subset \mathcal{T}^0$.

The names of these t-structures are explained by the following our main results in this subsection.

Theorem 4.10. (a) *We have mutually inverse bijections*

$$\mathcal{M} \mapsto (\mathcal{T}^{\leq 0}_{\mathcal{M}}, \mathcal{T}^{\geq 0}_{\mathcal{M}}) \quad \textit{and} \quad (\mathcal{T}^{\leq 0}, \mathcal{T}^{\geq 0}) \mapsto \mathcal{M}(\mathcal{T}^{\leq 0}, \mathcal{T}^{\geq 0})$$

between silting subcategories of $\mathcal{T}$ and silting t-structures of $\mathcal{T}$.
(b) *These induce bijections between tilting subcategories of $\mathcal{T}$ and tilting t-structures of $\mathcal{T}$.*

We need the following observation.

Lemma 4.11. *Let $(\mathcal{T}^{\leq 0}, \mathcal{T}^{\geq 0})$ be an arbitrary t-structure in $\mathcal{T}$ and $\mathcal{M} := \mathcal{M}(\mathcal{T}^{\leq 0}, \mathcal{T}^{\geq 0})$. Then*

$$\mathrm{Hom}_{\mathcal{T}}(\mathcal{M}, \mathcal{T}^{\leq 0}[>0]) = 0, \tag{16}$$

$$\mathrm{Hom}_{\mathcal{T}}(\mathcal{M}, \mathcal{T}^{\geq 0}[<0]) = 0, \tag{17}$$

$$\mathrm{Hom}_{\mathcal{T}}(\mathcal{M}, \mathcal{M}[>0]) = 0. \tag{18}$$

Proof. By $\mathcal{M} \subset {}^{\perp}(\mathcal{T}^{<0})$, we have (16). By $\mathcal{M} \subset \mathcal{T}^{\leq 0}$, we have (17). By $\mathcal{M} \subset \mathcal{T}^{\leq 0}$ and (16), we have (18). □

Now we are ready to prove Theorem 4.10.

(a)(i) Let $(\mathcal{T}^{\leq 0}, \mathcal{T}^{\geq 0})$ be a silting t-structure and $\mathcal{M} := \mathcal{M}(\mathcal{T}^{\leq 0}, \mathcal{T}^{\geq 0})$. Then $\mathcal{M}$ is skeletally small compact and generating by definition. Thus $\mathcal{M}$ is a silting subcategory by (18).

We have $\mathcal{T}^{\geq 0} \subset \mathcal{T}^{\geq 0}_{\mathcal{M}}$ by (17), and we have $\mathcal{T}^{\leq 0} \subset \mathcal{T}^{\leq 0}_{\mathcal{M}}$ by (16). Moreover we have

$$\mathcal{T}^{\leq 0} = {}^{\perp}(\mathcal{T}^{>0}) \supset {}^{\perp}(\mathcal{T}^{>0}_{\mathcal{M}}) = \mathcal{T}^{\leq 0}_{\mathcal{M}},$$
$$\mathcal{T}^{\geq 0} = (\mathcal{T}^{<0})^{\perp} \supset (\mathcal{T}^{<0}_{\mathcal{M}})^{\perp} = \mathcal{T}^{\geq 0}_{\mathcal{M}}$$

by Corollary 4.7, so $(\mathcal{T}^{\leq 0}, \mathcal{T}^{\geq 0}) = (\mathcal{T}^{\leq 0}_{\mathcal{M}}, \mathcal{T}^{\geq 0}_{\mathcal{M}})$ holds.

(ii) Let $\mathcal{M}$ be a silting subcategory of $\mathcal{T}$. Then $(\mathcal{T}^{\leq 0}_{\mathcal{M}}, \mathcal{T}^{\geq 0}_{\mathcal{M}})$ is a t-structure of $\mathcal{T}$ by Corollary 4.7. Let $\mathcal{N} := \mathcal{M}(\mathcal{T}^{\leq 0}_{\mathcal{M}}, \mathcal{T}^{\geq 0}_{\mathcal{M}})$. Then $\mathcal{N}$ is skeletally small since $\mathcal{T}^{\mathrm{c}}$ is skeletally small by Remark 2.9. We have

$$\mathrm{Hom}_{\mathcal{T}}(\mathcal{N}, \mathcal{N}[>0]) = 0 \tag{19}$$

by (18). Since $\mathcal{M}$ is silting, we have $\mathcal{M} \subset \mathcal{M}[<0]^{\perp} = \mathcal{T}^{\leq 0}_{\mathcal{M}}$. Since $\mathcal{T}^{<0}_{\mathcal{M}} = \mathcal{M}[\leq 0]^{\perp}$, we have $\mathrm{Hom}_{\mathcal{T}}(\mathcal{M}, \mathcal{T}^{<0}_{\mathcal{M}}) = 0$ and so $\mathcal{M} \subset {}^{\perp}(\mathcal{T}^{<0}_{\mathcal{M}})$. Consequently we have

$$\mathcal{M} \subset \mathcal{T}^{\leq 0}_{\mathcal{M}} \cap {}^{\perp}(\mathcal{T}^{<0}_{\mathcal{M}}) \cap \mathcal{T}^{\mathrm{c}} = \mathcal{N}. \tag{20}$$

Since $\mathcal{M}$ is generating, so is $\mathcal{N}$. By (19), we have that $\mathcal{N}$ is a silting subcategory of $\mathcal{T}$. In particular, $(\mathcal{T}^{\leq 0}_{\mathcal{M}}, \mathcal{T}^{\geq 0}_{\mathcal{M}})$ is a silting t-structure.

Now $\mathcal{M}$ and $\mathcal{N}$ are silting subcategories of $\mathcal{T}^{\mathrm{c}}$ in the sense of Definition 2.1 by Proposition 4.2. By (20) and Theorem 2.18 we have $\mathcal{M} = \mathcal{N}$. Thus our correspondences are mutually inverse.

(b) Let $\mathcal{M}$ be a tilting subcategory of $\mathcal{T}$. Then $\mathcal{M}$ is contained in both $\mathcal{T}^{\leq 0}_{\mathcal{M}}$ and $\mathcal{T}^{\geq 0}_{\mathcal{M}}$, so we have $\mathcal{M} \subset \mathcal{T}^{0}_{\mathcal{M}}$. Since $\mathcal{M}(\mathcal{T}^{\leq 0}_{\mathcal{M}}, \mathcal{T}^{\geq 0}_{\mathcal{M}})$ equals to $\mathcal{M}$, we have that $(\mathcal{T}^{\leq 0}_{\mathcal{M}}, \mathcal{T}^{\geq 0}_{\mathcal{M}})$ is a tilting t-structure.

Let $(\mathcal{T}^{\leq 0}, \mathcal{T}^{\geq 0})$ be a tilting t-structure and $\mathcal{M} := \mathcal{M}(\mathcal{T}^{\leq 0}, \mathcal{T}^{\geq 0})$. Since $\mathcal{M} \subset \mathcal{T}^{0}$, we have

$$\mathrm{Hom}_{\mathcal{T}}(\mathcal{M}, \mathcal{M}[<0]) \subset \mathrm{Hom}_{\mathcal{T}}(\mathcal{T}^{\leq 0}, \mathcal{T}^{>0}) = 0.$$

Thus $\mathcal{M}$ is a tilting subcategory of $\mathcal{T}$. □

References

[A] T. Aihara, *Silting mutation for self-injective algebras*, arXiv:1012.3265.

[AGI] T. Aihara, J. Grant, O. Iyama, in preparation.

[AST] I. Assem, M. Souto Salorio, S. Trepode, *Ext-projectives in suspended subcategories*, J. Pure Appl. Algebra 212 (2008), no. 2, 423–434.

[ASS] I. Assem, D. Simson, A. Skowronski, *Elements of the representation theory of associative algebras. Vol. 1. Techniques of representation theory*, London Mathematical Society Student Texts, 65. Cambridge University Press, Cambridge, 2006.

[APR] M. Auslander, M. I. Platzeck, I. Reiten, *Coxeter functors without diagrams*, Trans. Amer. Math. Soc. 250 (1979), 1–46.

[ARS] M. Auslander, I. Reiten, S. O. Smalo, *Representation theory of Artin algebras*, Cambridge Studies in Advanced Mathematics, 36. Cambridge University Press, Cambridge, 1997.

[BR] A. Beligiannis, I. Reiten, *Homological and homotopical aspects of torsion theories*, Mem. Amer. Math. Soc. 188 (2007), no. 883,

[BBD] A.A. Beilinson, J. Bernstein, P. Deligne, *Faisceaux pervers*, Analysis and topology on singular spaces, I (Luminy, 1981), 5–171, Asterisque, 100, Soc. Math. France, Paris, 1982.

[BGP] I. N. Bernstein, I. M. Gelfand, V. A. Ponomarev, *Coxeter functors, and Gabriel's theorem.* Uspehi Mat. Nauk 28 (1973), no. 2(170), 19–33.

[BB] S. Brenner, M. C. R. Butler, *Generalizations of the Bernšteĭn Gel'fand Ponomarev reflection functors*. Representation theory, II , pp. 103–169, Lecture Notes in Math., 832, Springer, Berlin-New York, 1980.
[BIRS] A. B. Buan, O. Iyama, I. Reiten, J. Scott, *Cluster structures for 2-Calabi-Yau categories and unipotent groups*, Compos. Math. 145 (2009), no. 4, 1035–1079.
[BMRRT] A. B. Buan, R. Marsh, M. Reineke, I. Reiten, G. Todorov, *Tilting theory and cluster combinatorics*, Adv. Math. 204 (2006), no. 2, 572–618.
[BRT] A. B. Buan, I. Reiten, H. Thomas, *Three kinds of mutation*, arXiv:1005.0276.
[BIKR] I. Burban, O. Iyama, B. Keller, I. Reiten, *Cluster tilting for one-dimensional hypersurface singularities*, Adv. Math. 217 (2008), no. 6, 2443–2484.
[C] W. Crawley-Boevey, *Exceptional sequences of representations of quivers*, Representations of algebras (Ottawa, ON, 1992), 117–124, CMS Conf. Proc., 14, Amer. Math. Soc., Providence, RI, 1993.
[GR] A. L. Gorodentsev, A. N. Rudakov, *Exceptional vector bundles on projective spaces*, Duke Math. J. 54 (1987), no. 1, 115–130.
[H] D. Happel, *Triangulated categories in the representation theory of finite-dimensional algebras*, London Mathematical Society Lecture Note Series, 119. Cambridge University Press, Cambridge, 1988.
[HU] D. Happel, L. Unger, *On a partial order of tilting modules*. Algebr. Represent. Theory 8 (2005), no. 2, 147–156.
[HK] M. Hoshino, Y. Kato, *Tilting complexes defined by idempotents*, Comm. Algebra 30 (2002), no. 1, 83–100.
[HKM] M. Hoshino, Y. Kato, J. Miyachi, *On t-structures and torsion theories induced by compact objects*, J. Pure Appl. Algebra 167 (2002), no. 1, 15–35.
[HX] W. Hu, C, Xi, *D-split sequences and derived equivalences*, Adv. Math. 227 (2011), no. 1, 292–318.
[IO] O. Iyama, S. Oppermann, *n-representation-finite algebras and n-APR tilting*, arXiv:0909.0593, to appear in Trans. Amer. Math. Soc.
[IR] O. Iyama, I. Reiten, *Fomin-Zelevinsky mutation and tilting modules over Calabi-Yau algebras*, Amer. J. Math. 130 (2008), no. 4, 1087–1149.
[IW] O. Iyama, M. Wemyss, *Auslander-Reiten duality and maximal modifications for non-isolated singularities*, arXiv:1007.1296.
[IY] O. Iyama, Y. Yoshino, *Mutation in triangulated categories and rigid Cohen-Macaulay modules*, Invent. Math. 172 (2008), no. 1, 117–168.
[K] B. Keller, *Deriving DG categories*, Ann. Sci. Ecole Norm. Sup. (4) 27 (1994), no. 1, 63–102.
[KV] B. Keller, D. Vossieck, *Aisles in derived categories*, Deuxieme Contact Franco-Belge en Algebre (Faulx-les-Tombes, 1987). Bull. Soc. Math. Belg. Ser. A 40 (1988), no. 2, 239–253.
[KY] , B. Keller, D. Yang, *Derived equivalences from mutations of quivers with potential*, Adv. Math. 226 (2011), no. 3, 2118–2168.
[KM] D. Kussin, H. Meltzer, *The braid group action for exceptional curves*, Arch. Math. (Basel) 79 (2002), no. 5, 335–344.
[L] S. Ladkani, *Perverse equivalences, BB-tilting, mutations and applications*, arXiv:1001.4765.
[M] J. Miyachi, *Localization of triangulated categories and derived categories*, J. Algebra 141 (1991), no. 2, 463–483.
[N] A. Neeman, *The connection between the K-theory localization theorem of Thomason, Trobaugh and Yao and the smashing subcategories of Bousfield and Ravenel*, Ann. Sci. Ecole Norm. Sup. (4) 25 (1992), no. 5, 547–566.
[O] T. Okuyama, *Some examples of derived equivalent blocks of finite groups*. preprint, 1998.
[P1] D. Pauksztello, *Compact corigid objects in triangulated categories and co-t-structures*, Cent. Eur. J. Math. 6 (2008), no. 1, 25–42.
[P2] D. Pauksztello, *A note on compactly generated co-t-structures*, arXiv:1006.5347.
[Ric1] J. Rickard, *Morita theory for derived categories*, J. London Math. Soc. (2) 39 (1989), no. 3, 436–456.
[Ric2] J. Rickard, *Derived categories and stable equivalence*. J. Pure. Appl. 61 (1989), 303–317.
[RS] C. Riedtmann, A. Schofield, *On a simplicial complex associated with tilting modules*, Comment. Math. Helv. 66 (1991), no. 1, 70–78.
[Rin] C. M. Ringel, *The braid group action on the set of exceptional sequences of a hereditary Artin algebra*, Abelian group theory and related topics (Oberwolfach, 1993), 339–352, Contemp. Math., 171, Amer. Math. Soc., Providence, RI, 1994.
[SY] Y. Sekiya, K. Yamaura, *Tilting theoretical approach to moduli spaces over preprojective algebras*, arXiv:1008.4451.
[W] J. Wei, *Semi-tilting complexes*, preprint.

T. Aihara: Division of Mathematical Science and Physics, Graduate School of Science and Technology, Chiba University, Yayoi-cho, Chiba 263-8522, Japan
E-mail address: taihara@math.s.chiba-u.ac.jp

O. Iyama: Graduate School of Mathematics, Nagoya University, Chikusa-ku, Nagoya, 464-8602 Japan
E-mail address: iyama@math.nagoya-u.ac.jp